
ANN. REP WLM 28 .BE 5 B 84 1961-63

THE BETHLEM ROYAL HOSPITAL AND THE MAUDSLEY HOSPITAL



TRIENNIAL STATISTICAL REPORT

YEARS 1961-1963


Edited by E. H. HARE



THE BETHLEM ROYAL HOSPITAL AND THE MAUDSLEY HOSPITAL

TRIENNIAL STATISTICAL REPORT

YEARS 1961-1963



Edited by
E. H. HARE, M.A., M.D., D.P.M.

Published by the Bethlem Royal Hospital and the Maudsley Hospital, Denmark Hill, London 1965

WELLCOME							
Ann Rep							
WLM 28							
8 July Some Supple							
B & 4							
1961-63							

FOREWORD

The present Report is somewhat bulkier than its two predecessors. This is because it not only presents the statistics of the current triennium but compares them with those of the four previous triennia and discusses some of the statistical trends discernable over this period. As Reports accumulate, the study of such trends should be of increasing historical interest; it may also perhaps have a practical value, though how far a trend reflects psychiatric progress rather than a change in fashion, selection, distribution or natural history, will never be easy to determine. With the hope of emphasising the historical interest of hospital statistics, I have made a brief comparison of the present triennial figures with those of Bethlem Hospital a hundred years ago (Chapter I).

A change in lay-out from previous reports may be noted. Percentages are given to the nearest whole number and not, as formerly, to a decimal place. To help distinguish numbers from percentages, all percentage figures in the tables are printed in italics.

Professor Sir Aubrey Lewis and Dr. Michael Shepherd read the manuscript, and I am grateful to them for their criticisms.

E. H. HARE
The Maudsley Hospital
Denmark Hill
London, S.E.5

February, 1965

COMMENTARY

						p	age
Foreword	• • •	• • •	• • •	• • •	• • •	• • •	iii
Chapter One. INTRODUCTION							
1 T D		• • •	• • •	• • •			1
2. Summary of Main Trends D				• • •	• • •	• • •	1
2 A 71 37- A				• • •	• • •	• • •	2
4. New Features of the Presen			• • •			• • •	5
5 Decision of Transco		OKI	• • •	• • •	• • •	• • •	5
3. DEFINITION OF TERMS	• • •	• • •	• • •	• • •	• • •	• • •	3
Chapter Two. ADULTS: SOCIAL	DATA	Λ.					
1. Numbers of Patients .		• • •		• • •		• • •	8
2. Age and Sex		• • •	• • •		• • •	• • •	8
3. Previous Discharges			• • •		• • •	• • •	8
4. Religion		• • •	• • •			• • •	9
5. OCCUPATION AND SOCIAL CLA	ASS	• • •	• • •			• • •	9
6. Marital Status			• • •			• • •	9
7. Patients Who Asked to be l	Refere	RED			• • •	• • •	9
8. Country of Birth		• • •		• • •	• • •	• • •	9
Chapter Three. ADULTS: HOSPIT	AL D	ATA					
1. Referring Agencies .		• • •		• • •		• • •	19
2. Duration of Stay		• • •	• • •			• • •	19
3. Number of Attendances		• • •	• • •	• • •	• • •	• • •	20
4. Special Investigations		• • •	• • •	• • •		• • •	20
5. Special Treatment (In-patie	ENTS)	• • •	• • •			• • •	20
6. OUTCOME OF TREATMENT (IN-	PATIEN	TS)	• • •		• • •	• • •	21
7. Disposal			• • •			• • •	21
8. Lapses in Treatment .		• • •	• • •		• • •	• • •	21
9. Travelling Time to Hospita	AL		• • •				22
Chapter Four. ADULTS: DIAGNO	OSTIC	DATA	A				
1. Numbers of Patients with V	ARIOU	s Diac	NOSES	• • •	• • •	• • •	32
2. Miscellaneoùs Diagnoses .		• • •	• • •	• • •		• • •	33
3. Asked to be Referred .		• • •		• • •	• • •	• • •	34
4. Lapses in Treatment .	• • •	• • •	• • •	• • •		• • •	34
Chapter Five. CHILDREN							
A. THE ADOLESCENT UNIT BY P. H. CONNELL AND		 RREN	• • •	• • •	• • •	•••	40
B. Commentary on the Table	S		• • •	• • •	• • •	• • •	42
C. Brixton Child Guidance (CLINIC		• • •	• • •	• • •	• • •	42

Chapter Six.		
A. DAY-PATIENTS	• • •	59
B. DOMICILIARY VISITS	• • •	59
Chapter Seven. SPECIAL CLINICS		
A. The Out-patient Department and Emergency Clinic		
1. HISTORICAL REVIEW BY D. H. BENNETT	• • •	66
2. Notes on the Tables	• • •	68
3. Out-patient Waiting Times	• • •	68
B. The Psychiatric Emergency Unit, St. Francis' Hospital		69
C. The Forensic Unit	• • •	69
D. The Guy's-Maudsley Neurosurgical Unit		70
Chapter Eight. GENERAL PRACTITIONER REFERRALS		77
APPENDIX		
Front Page for Adults and Children		82
CODING SHEET FOR DOMICILIARY SERVICE	• • •	86

TABLES

Cha	pter (One	p	age
	1.1	Numbers of new patients, patients, and discharges, 1961-63		X
	1.2	Average number of in-patient beds available		7
	1.3	Numbers of hospital professional staff	• • •	7
Cha	pter 7	Two. Adults		
	2.1	Numbers of adult patients and discharges in five triennia		10
	2.2	In-patient discharges, by hospital and year		10
	2.3	Age and sex of in-patients and out-patients		11
	2.4	Previous in-patient discharges (before 1961) of in-patients		11
	2.5	Previous out-patient discharges (before 1961) of out-patients		12
	2.6	Religion		12
	2.7	Occupation		13
	2.8	Social class of males for five triennia		14
	2.9	Marital status		14
	2.10	Number of children born to hospital patients		15
	2.11	Birth order	• • •	15
	2.12	Twins, etc	• • •	16
	2.13	Asked to be referred		16
	2.14	Previous treatment at other hospitals		17
	2.15	Relatives treated		17
	2.16	Country of birth	• • •	18
	2.17	Country of birth: hospital patients and London populations		18
Cha	ipter T	Three. Adults		
	3.1	Referring agencies for in-patients		22
	3.2	Referring agencies for out-patients		23
	3.3	Referring agencies to the Out-patient Department	• • •	23
	3.4	Self-referrals, by social class		24
	3.5	Median and average duration of in-patient stay for five trienn	ia	24
	3.6	Duration of in-patient stay		24
	3.7	Duration of in-patient stay, by social class	• • •	25
	3.8	Number of out-patient attendances		25
	3.9	Special investigations	• • •	26
	3.10	Comparison of special treatments for five triennia (in-patients))	26
	3.11	Special treatments (in-patients)	• • •	27
	3.12	Outcome of treatment (in-patients)		27
	3.13	Outcome of treatment, by social class	• • •	27
	3.14	Causes of death in in-patients	• • •	28
	3.15	Disposal of in-patient discharges	• • •	28
	3.16	Disposal of out-patient discharges	• • •	29
	3.17	Lapses in treatment	• • •	29

							1	vage
	Lapses in treatment, by soc			• • •	• • •	• • •	• • •	30
3.19	Travelling time from home	to hos	spital (out-par	tient a	ittendar	nces)	
	Travelling time, by social c			• • •	• • •	• • •	• • •	30
	Travelling time, by number		ndance	es	• • •	• • •	• • •	31
3.22	Travelling time, by disposal	l, etc.	• • •	• • •	• • •	• • •	• • •	3
Chapter	Four. Adults							
4.1	Diagnoses in four major gro	oups	• • •	• • •	• • •	• • •	• • •	34
4.2	Diagnoses in four major gro	oups, fo	or five	triennia	ı	• • •		35
4.3	Diagnoses of in-patients	• • •	• • •	• • •				35
4.4	Diagnoses of in-patients, fo	r five tr	iennia				• • •	36
4.5	Diagnoses of out-patients	• • •	• • •	• • •				36
4.6	Diagnosis of out-patients,	for five	trienn	nia	• • •			37
4.7	Miscellaneous diagnoses.—				• • •			37
4.8	Principal accessory diagnose				S			38
4.9	Marital status, by diagnosis		-	_		• • •		38
4.10								39
4.11	Lapses in treatment, by diag		···					39
	1	5				•••	•••	
Chapter 1	Five. CHILDREN							
5.1	Numbers of patients and dis	charge	for f	ive tries	nio			45
5.2	In-patient discharges, by hos					• • •	• • •	45
5.3	A	spitai ai	nu yea.	1	• • •	• • •	• • •	
	Religious upbringing	• • •	• • •	• • •	• • •	• • •	• • •	46
5.5			* * *	• • •	• • •	• • •		46
	Social class of parents		• • •	• • •	• • •	• • •	• • •	47
5.6	Social class of parents, for f		ınıa		• • •	• • •	• • •	47
5.7	Marital status of mother	• • •	• • •	• • •	• • •	• • •	• • •	47
5.8	Mother's age at marriage		• • •	• • •	• • •	• • •	• • •	48
5.9	Mother's age at patient's bir		• • •	• • •	• • •	• • •	• • •	48
	Sibship size		• • •	• • •	• • •	• • •	• • •	49
	Birth order, by sibship size			• • •	• • •	• • •	• • •	49
	Twins, etc		• • •	• • •	• • •	• • •	• • •	50
5.13	Cared for by foster parents,	etc.	• • •		• • •		• • •	50
5.14	Relatives treated		• • •	• • •		• • •	• • •	51
5.15	Referring agencies for in-pat	tients	• • •	• • •	• • •	• • •	• • •	51
5.16	Referring agencies for out-pa	atients	• • •	• • •	• • •	• • •	• • •	52
5.17	Duration of in-patient stay		• • •	• • •		• • •	• • •	52
5.18	Median duration of in-patien	nt stay,	for fiv	e trienr	nia	• • •		53
	Number of out-patient atten			• • •	• • •			53
5.20	Number of out-patient atten	dances,	for fiv	e trieni	nia	• • •		53
	Special investigations.—In-pa			• • •	• • •	• • •		54
	Special investigations.—Out-		8	• • •	• • •	• • •		54
	Special treatments					• • •		55

								p	age
5.24	Outcome of treatment		• • • •	• •	• • •		• • •	• • •	55
5.25	Outcome of treatment	, for fiv	ve trien	nia		• • •	• • •	• • •	56
5.26	Disposal	• • •		• •		• • •	• • •	• • •	56
5.27	Lapses in treatment	• • •	• • •		• • •	• • •	• • •	• • •	57
5.28	Lapses in treatment, b	y age	• • •		• • •	• • •	• • •	• • •	57
5.29	Diagnosis	• • •		• • •		• • •		• • •	57
5.30	Country of birth of ch	nildren'	s paren	ts		• • •	• • •	• • •	58
5.31	Staffing at Brixton Ch	nild Gu	idance	Clinic		• • •	• • •	• • •	58
5.32	Case statistics at Brix	ton Chi	ild Gui	dance	Clinic	• • •	• • •	• • •	58
Chantay S	Tina.								
Chapter S									
	Y-PATIENTS		1	, • •					(0
6.1	Number of discharges	s, by ye	ar and	trienni	a	• • •	• • •	• • •	60
6.2	Age	• • •	• • •	• • •	• • •	• • •	• • •	• • •	60
6.3	Social class	• • •	• • •	• • •	• • •	• • •	• • •	• • •	61
6.4	Marital status	• • •	• • •	• • •	• • •	• • •	• • •	• • •	61
6.5	Referring agency		• • •	• • •	• • •	• • •	• • •	• • •	61
6.6	Duration of stay		• • •	• • •	• • •	• • •	• • •	• • •	62
6.7	Special treatments		• • •	• • •	• • •	• • •	• • •	• • •	62
6.8	Outcome of treatmen	t	• • •	• • •	• • •	• • •	• • •	• • •	63
6.9	Diagnosis	• • •	• • •	• • •	• • •	• • •	• • •	• • •	63
B. Do	MICILIARY PATIENTS								
6.10	Domiciliary visits for	five tri	ennia		• • •	• • •	• • •		63
	Age		• • •	• • •		• • •			64
	Marital status	• • •	• • •	• • •	• • •	• • •			64
6.13	Social class	• • •	• • •	• • •	• • •	• • •		• • •	64
6.14	Recommended dispos	sal	• • •	• • •					65
	Diagnosis		• • •	• • •	• • •	• • •	• • •	• • •	65
Chapter .	Seven								
A. Em	ERGENCY CLINIC								
7.1	New patients referred	l to the	Out-pa	itients,	by trie	ennia	• • •	• • •	70
7.2	Emergency clinic atte	endance	es, by ye	ear	• • •	• • •	• • •	• • •	70
7.3	Numbers, by year	• • •	• • •	• • •	• • •		• • •	• • •	70
7.4	Referring agencies	• • •	• • •	• • •		• • •	• • •	• • •	70
7.5	New patients, by age		•••	• • •		• • •	• • •	• • •	71
7.6	Diagnosis of new pat	tients	• • •	• • •	• • •	• • •	• • •	• • •	71
Out	-PATIENT WAITING TIM	MES							
7.7	Out-patient waiting t	imes, b	y year	• • •		• • •			72
7.8	Out-patient waiting t	· ·							72

								ρ	uge
B. EM	MERGENCY UNIT (St. FI	rancis	' Hospi	tal)					
7.9	Status on admission	• • •	• • •			• • •	• • •		72
7.10) Age	• • •				• • •			73
7.11	Marital status	• • •							73
7.12	2 Diagnosis						• • •		73
7.13	Referring agency								74
7.14	4 Mode of discharge			• • •	• • •	• • •	• • •		74
C. Fo	DRENSIC UNIT								
7.13	5 Numbers of new cas	ses and	d report	is		• • •			75
D. N	eurosurgical Unit								
	6 Numbers of patients	and o	operatio	ons				• • •	75
	7 Leucotomies and op		_		y	• • •		• • •	75
	8 Attendances at X-ra				• • •	• • •			76
Chapter	Fight								
8.1	Number of general	pract	itioners	and c	of new	cases 1	referred	l. bv	
0.1					• • •	•••		•••	80
8.2	1			ith Lo	ndon p	ostal	districts	s, by	-
	number of new cas			• • •	• • •	•••		•••	81
8.3	Number of new car London areas	ses re	ferred :	per pra	actition 	er, for	five S	outh	81
Map									
-	w cases, referred by	gener	al prac	ctitione	ers, fro	m Lor	ndon p	ostal	
	listricts	• • • •	• • •		* * *	• • •	• • •		3-79

Table 1.1 Numbers of new patients, patients, and discharges¹, 1961-63

			0	
		Male	Female	Total
ADULTS				
New Patients				
Hospital patients		4,184	4,280	8,464
In-patients	• • • • • •	1,413	1,913	3,326
Out-patients	•••	3,112	2,782	5,894
Patients				
Hospital patients	• • • • • • •	5,539	5,963	11,502
In-patients	* * *	1,676	2,272	3,948
Out-patients	•••	3,960	3,806	7,766
Discharges				
Total	• • • • • • • • • • • • • • • • • • • •	6,217	6,809	13,026
In-patient	•••	1,920	2,689	4,609
Out-patient	•••	4,297	4,120	8,417
CHILDREN				
New patients				
Hospital patients	• • • • • •	650	405	1,055
In-patients	• • • • • • • • • • • • • • • • • • • •	174	134	310
Out-patients	•••	483	274	757
Patients				
Hospital patients	• • • • • • • •	683	415	1,098
In-patients		180	134	314
Out-patients	• • • • • • • • • • • • • • • • • • • •	503	281	784
Discharges				
Total	• • • • • • •	690	421	1,111
In-patient	• • • • • •	185	138	323
Out-patient	• • •	505	283	788

¹See definitions, Chapter 1

CHAPTER ONE

INTRODUCTION

1. THE FIFTH REPORT

The present triennial statistical report, the fifth in the series, deals with cases discharged during the years 1961-63 and also with a comparison of statistics over five triennia. The purpose, method of collection, and essential limitations of these statistics have been described and discussed in earlier reports. It may be repeated, however, that the reports contain the only detailed statistics on psychiatric out-patients so far published in Great Britain.

2. TRENDS DURING THE PAST 15 YEARS

The first five Statistical Reports of the Bethlem Hospital cover the years 1949-1963. Statistical trends over this 15-year period can be discerned in the appropriate tables and are discussed in the commentaries, but for ease of reference the most interesting and clear-cut of these trends are summarised below. It must be remembered that the existence of a statistical trend tells us nothing about its cause and that the causal explanations offered here are no more than likely possibilities.

- (1) Numbers of Patients. There has been a steady and marked increase in the numbers of adult patients attending the hospital. The over-all increase in 15 years has been 50% (p. 8). For outpatients, the rate of increase has been growing since 1952, and this is probably due in part to the increasing use being made of the Emergency Clinic. There has been no increase in the numbers of children; the numbers of out-patient children are falling (p. 42).
- (2) Duration of Stay. The duration of stay of adult in-patients has steadily decreased (pp. 19, 24), such that over the last twelve years there has been a 30% reduction for both sexes in the time of stay. For children, the median duration of stay has decreased for girls, but for boys it has increased (p. 44).
- (3) Special Investigations (pp. 20,26). X-ray examination of adult patients has become progressively more common, the proportion of cases X-rayed increasing from 23% to 55%. The use of blood counts and bacteriological tests has also increased, but that of electroencephalography, basal metabolic rate and special psychological tests of intelligence has decreased. The decrease in psychological tests probably reflects the fact that simple intelligence tests are increasingly administered by the medical staff.
- (4) Special Treatments (pp. 20,27). Coma insulin therapy declined after 1952-54 and was discontinued altogether in 1958. Modified insulin therapy has progressively declined from 65 cases per year to 11 per year. The number of leucotomy operations, which at first rose from 11 per year to 30 per year (in 1952-54),

has since fallen again to 10 per year. The proportion of cases receiving electro-convulsive therapy rose between 1949 and 1957 from 29% to 34%, but has since fallen back to 25%. It is thus clear that what have been called the physical treatments of psychiatry are being used less. Their decline has been off-set by an increased use of drugs.

- (5) Diagnosis (p. 32). Among adult in-patients there has occurred a steady and notable decrease in the proportion of cases diagnosed as neurotic. This is due to a decrease in anxiety neurosis, hysteria and obsessional neurosis, but not in neurotic depression. The proportion of cases diagnosed as manic-depressive psychosis has increased greatly (from 18% to 31%). Psychopathic disorder and alcoholism have also increased, but only since 1958. Similar trends occur in the diagnoses of out-patients (p. 36) and day-patients (p. 63). Possible causes are discussed in Chapter IV.
- (6) Mode of Discharge. There has been a progressive decrease, from 45 per year to 9 per year, in the number of cases where transfer from the hospital to an Observation Ward (or Emergency Unit) was necessary (p. 21). Every such transfer represents a failure on the part of the hospital to continue treatment on a voluntary or informal basis, and the fall in numbers may reflect an increased efficacy of drug treatment in the management of disturbed cases.

The proportion of in-patients who discharged themselves against medical advice decreased slightly (from 14% to 12%) during fifteen years. But there was an increase, and a fairly marked one (from 19% to 28%), in the proportion of out-patients lapsing in attendance (p.21). Among out-patient children a similar trend is apparent (p. 44).

(7) Outcome of Treatment. The proportion of adult in-patients discharged as recovered or much improved has increased steadily (from 47% to 53%), with a corresponding decrease in the proportion discharged as unchanged or worse (p. 21). Among inpatient children, however, trends opposite to these occurred (p. 44). It would be hazardous to guess the probable causes of these trends.

To summarise in one sentence the main trends of the past fifteen years: the number of adult patients coming to the hospital is increasing; the duration of stay of in-patients is decreasing; physical methods of treatment are giving place to drugs; anxiety neurosis is being diagnosed less and manic-depressive psychosis more; fewer patients are needing to be transferred on account of disturbed behaviour; and a small but noteworthy increase is occurring in the proportion of patients discharged as recovered or much improved.

3. A HUNDRED YEARS AGO

Hospital statistical reports provide the historian with a record not only of numbers and types of cases but also of the varieties of information thought at the time to be worth collecting; and what is thought worth collecting will depend not only on the technical development of the subject but also on prevailing ideas about causation and prognosis. Much of the present report is concerned with a comparison of figures in the past five triennia, and it may therefore be of interest to devote a page or two to the Bethlem

statistics of a hundred years ago.

During 1861-63, Bethlem hospital was in St. George's Fields, Lambeth, on the site of what is now the Imperial War Museum. The hospital cases at that time were classed as curable, incurable, and criminal. The criminal cases (numbering about 130) were transferred to Broadmoor in 1863-64; the incurable cases (numbering about 70) were those not fit for discharge after at least a year's stay in the curable section and for whom a bed was available; the curable cases were cases thought to be of good prognosis and admitted under the provision that they would not ordinarily stay more than a year. It is the admission and discharge of the curable cases which were the main statistical concern of the Resident Physician of Bethlem in his annual reports and it is these cases which are fairly comparable with those nowadays admitted to the Bethlem-Maudsley hospital.

The average annual number of discharges during 1861-63 was 60 male and 101 female cases. Their median age (31 years for males, 35 for females) was five years younger than in 1961-63, and 45% were unmarried compared with 36% now. Acute melancholia was the commonest diagnosis, made in 30%. No category of chronic melancholia is listed and the next commonest diagnosis was chronic mania (22%), followed by acute mania (18%) and dementia (18%). Puerperal mania was recorded in 7% of females¹. Cases of general paralysis were not admitted to Bethlem because such cases were incurable; but doubtful cases were given the benefit, and the increasing proportion of admissions (5% in 1861-63) in which this diagnosis was eventually made led Sir Charles Hood to lament "how manifestly a form of mental disease known as general paralysis appears to be on the increase".

In each annual report a table lists the total of curable patients admitted during the previous hundred years. For the centennium 1762-1861 this total was 19,624 of whom 44% were discharged cured and 7% died in hospital. For 1861-63 the proportion discharged cured was 55%, a figure which may be compared with the 53% discharged recovered or much improved during 1961-63.²

¹ And by one of those slips from which no statistical report can claim immunity, this diagnosis is also assigned to 7 males. See also *Brit. Med. J.*, 1965, i, 401, for a grosser example and some pertinent comment.

² Fluctuations in the cure rate might be the subject of an interesting study. When Edward Tyson was physician to Bethlem (from 1684 to 1703), 69 per cent of the patients were "cured and discharged". See Leigh, D., *The Historical Development of British Psychiatry* (London, 1961, page 2).

The proportion of deaths in 1861-63 was 8%, the commonest cause of death being exhaustion following acute mania or melancholia. In 1961-63, the death rate among an older population of in-patients was less than 1% and though administrative changes may provide a partial explanation yet this fall in death rate is perhaps the most striking of the differences to be remarked in a comparison of these centenary statistics.

Many statistics in the old reports have no counterpart now, and the reasons for this may variously be assigned to changes in frequency of the events, in ideas of the reliability of information, or simply in administrative concepts of what figures should be taken to reflect a progresive hospital. In 1861-63, 42% of the patients are recorded as having meditated or attempted suicide before admission, while 34% were at the time of admission classed as dangerous or violent to others. Today, only a minute proportion of Bethlem patients could be described as dangerous or violent; and even if we included the Psychiatric Emergency Unit at St. Francis' Hospital, the proportion would still be very small². Yet in spite of the high proportion of violent admissions, the weekly average of patients under restraint was nil and had been so since 1851. We learn from another table that of those discharged cured 41% had been ill for less than a month before admission, while of those discharged uncured this proportion was only 19%—a contrast which led the resident physician, William Helps, to observe that "it cannot be too strongly impressed ... how much a favourable result in the treatment of the insane depends on early and judicious management", and to regret that the loss of opportunity for early treatment "can only be accounted for by supposing that a false notion exists," that by becoming an inmate of an Asylum a person incurs an ineradicable social stain".

Aetiological factors in mental illness seemed less obscure a century ago than they do today. In a table of "apparent or assigned causes", puerperal or uterine disturbance is the commonest in females (20% of cases) and masturbation in males (12%). Heredity accounts for 10%, mental anxiety for 8%; then come loss of relatives (6%), reduced circumstances (5%), and intemperance, surprisingly low at 4%. Other causes of mental illness (leading to the social stain of asylum admission) include disappointment, fright, excessive study and religious excitement. "Previous attacks" was considered a sufficient cause and covers 15% of cases, while in a final 15% the cause is "not traceable, probably idiopathic". The degree of education, anticipating our present social class by occupation, was rated superior in 8%, good in 35%, moderate in 56% and indifferent or none in 1%; but no analysis of diagnosis, or of outcome, by education is attempted. The proportion of patients

² This and other differences between Bethlem patients a hundred years apart have been described and discussed by Klaff and Hamilton (*J. Ment. Sci.*, 1961, **107**, 819).

attending Sunday chapel in 1861 is recorded as 55.80%, a figure which declined during the triennium to 53.13%.

4. NEW FEATURES OF THE PRESENT REPORT

The work of the adolescent unit (at Bethlem) is described in Chapter V, and that of the Out-patients Department in Chapter VII. Chapter VII also gives statistics for discharges from the Psychiatric Emergency Unit at St. Francis' Hospital, Dulwich, which is staffed by doctors from the Bethlem-Maudsley hospital.

New tables deal with the patients' country of birth (Tables 2.17, 5.30), with self-referrals by social class (Table 3.3) and with an analysis of factors associated with the time taken by patients to travel from home to the Out-patient Department (Tables 3.19-3.22). Also new are the tables dealing with cases first seen at the Emergency Clinic (Tables 7.1-7.4) and with waiting-times for out-patient appointments (Tables 7.7, 7.8).

In the present report, percentage figures have been given in whole numbers and not (as previously) to a decimal place; in the

tables, percentages are printed in italics.

5. DEFINITION OF TERMS

The same definitions are adopted as in the previous two reports. They are as follows:

A. Adults and Children. Adult patients are defined as those admitted to the adult departments of the hospital; with very few exceptions, adult patients are aged 16 or over.

Patients described in the report as children are those admitted to the children's departments and are, with very few exceptions,

aged under 16 at the time of admission.

B. The Hospital and its Departments. The word "hospital" is here taken to cover the in-patient and out-patient departments

of the Royal Bethlem Hospital and the Maudsley Hospital.

For adults, the in-patient department includes wards at Bethlem and the Maudsley but does not include the wards of the Guy's Maudsley Neurosurgical Unit. The adult out-patient department includes the Maudsley out-patient department, the Emergency clinic, the day-hospitals at Bethlem and at the Maudsley, and all follow-up clinics.

For children, the *in patient department* includes the children's in-patient unit at the Maudsley Hospital and the adolescent ward at Bethlem; the *out patient department* is at the Maudsley Hospital.

¹ The figure continued to decline steadily, and in 1882 it was only 32.2 per cent. But we should resist the temptation to conclude that this was why it was omitted altogether from the reports after that year.

- C. Admissions and Spells of Care. A period of time during which a patient remains continuously under care at the hospital, without being discharged or lapsing in attendance, is called a spell of care. Each spell of care begins with the admission of the patient and ends with his discharge. The meaning of the term admission is limited by the hospital's "three-months rule"; the rule is that if a person comes under the care of the out-patient department within three months of being discharged from either of the departments, then this does not count as a new admission but is considered simply as a continuation of his previous spell of care.
- D. Discharge. An in-patient discharge is the discharge of a patient at the end of a spell of care which included a period of in-patient care.

An *out-patient discharge* is the discharge of a patient at the end of a spell of care which did *not* include a period of in-patient care.

A hospital discharge is the discharge of a patient at the end of any

spell of care.

Because a hospital discharge must be either an in-patient discharge or an out-patient discharge, the total number of hospital discharges equals the sum of the in-patients and out-patient discharges (see Table 1.1).

E. Patients Discharged. During any triennium, many patients have more than one spell of care at the hospital. For this reason, the number of individual patients discharged from a department or from the hospital is less than the number of discharges. Patients are classed as **in-patients**, **out-patients**, or **hospital patients** according to the type of discharge with which their spells of care are associated.

Because a patient may be discharged as an in-patient on one occasion and as an out-patient on another occasion, the sum of in-patients and out-patients will in general be greater than the number of hospital patients (see Table 1.1); but, to the first approximation, hospital patients may be thought of as the sum of in-patients and out-patients.

F. New Patients. These are patients who, during the triennium, attend and are discharged from the hospital for the first time in their lives. A new in-patient is one who completes his first-ever spell of in-patient care; a new out-patient is one who completes his first-ever spell of out-patient care; and a new hospital patient is one who completes his first-ever spell of care at the hospital, whether as an in-patient or as an out-patient.

Because a patient may qualify as a new in-patient on one occasion and as a new out-patient on another occasion, the sum of new inpatients and new out-patients will in general be greater than the

number of new hospital patients (Table 1.1).

G. Cases. The word case has been used loosely in the report. Its appropriate meaning is mostly obvious from the context, but in general it has been taken to refer to the illness of a patient receiving a particular spell of care.

Table 1.2 Average number of in-patient beds available during 1961-63

Department		Maudsley	Bethlem	Joint Hospital
Psychiatric Adults Children Neurosurgical	• • •	197 26 30	199 30 —	396 56 30
TOTAL	• • •	253	229	482

Table 1.3 Numbers of hospital professional staff

				, ,	
	1949	1954	1957	1960	1963
DOCTORS ²					
Consultants			2		
Whole-time	10	8	9	8	8
Part-time (and whole- time equivalent in	Î				
brackets)	8	14	15 (9)	18 (9)	30 (12)
Senior Registrars				` ,	
Whole-time	Į.				
Part-time Registrars and Senior House	44	60	67	64	65
Officers		00	07	0-1	03
Clinical Assistants	1				
NHIDGEG					
NURSES Whole-time					
Male	Ì			74	84
Female	182	247	237	173	227
Part-time	1			4	3
Male } Female }	49	84	107	92	82
1 cintaie y	1		107		0.2
CLINICAL PSYCHOLOGISTS	6	10	9	8	9
PSYCHIATRIC SOCIAL WORKERS	11	11	12	14	16
F ST CHIATRIC SOCIAL WORKERS	11	11	12	14	10
OCCUPATIONAL THERAPISTS	9	12	12	13	15

¹Excluding the Neurosurgical Unit

CHAPTER TWO

ADULTS: SOCIAL DATA

Introduction

This chapter deals with the demographic and social aspects of patients. The tables are based on the number of patients attending the hospital or its departments. For those patients who were discharged more than once during a triennium, the information used is that recorded at the first discharge.

1. Numbers of Patients

Table 2.1 shows the continued steady increase in the number of cases dealt with at the hospital during five triennia. In 1961-63 the numbers of in-patients, of out-patients, and of total discharges were all 50% higher than in 1949-51. Between 1955-57 and 1961-63 (the figures for earlier triennia were not extracted), the number of new in-patients increased by 14% and the number of new out-patients by 32%. It is thus clear that, in terms of care given to patients, the volume of work done in the hospital has greatly increased over the past fifteen years. Moreover, as there has been no significant increase in the number of in-patient beds or in the space of the out-patient department during this period, the increased work is reflected for in-patients in an increased turn-over of cases, and for out-patients in an increased number of clinics and number of cases per clinic.

Table 1.1 (page x) shows that during 1961-63 there were 8,464 new adult patients, i.e., patients discharged for the first time from the hospital The total discharges were 13,026 so that the redischarge rate was 35%. This rate is notably less than in the previous two triennia (49% and 43%); the diminution is largely to be accounted for by the increased numbers of new cases seen at the Emergency Clinic. For in-patients, the re-discharge rate was 28%, for outpatients, 30%.

2. Age and Sex (Table 2.3)

Since 1955-57 the proportion of hospital patients under 25 years old has increased from 15% to 20%, and the proportion of those over 55 has decreased from 17% to 11%. Thus the hospital population is at present tending to become younger.

3. Previous Discharges (Tables 2.4, 2.5)

There has been no important change during five triennia in the proportion of discharged patients who had also been discharged from the hospital in previous years. This proportion is about 15%.

Of in-patients discharged during 1961-63, 11 males and 16 females had previously been in-patients in the Children's Department and 21 males and 18 females had been out-patients there. For out-patients, these numbers were 17 males and 14 females, and 61 males and 38 females respectively.

4. Religion (Table 2.6)

Over five triennia, the proportion of patients who belonged to the Church of England has steadily decreased (from 74% to 65%). This has been balanced by an increase in the proportion of Roman Catholics (from 11% to 18%) and of those giving their religion as "none" (from 2% to 5%).

5. OCCUPATION AND SOCIAL CLASS (Tables 2.7, 2.8)

Among males, a noteworthy change in occupational rates from the previous triennium is the increase in the professional and technical group from 11% to 14% (i.e., from 527 patients to 756).

The number of patients who are students (at school, college or university) continues to increase. In successive triennia since 1952, the numbers have been: Males, 107, 180, 225, 280; Females, 58, 61, 88, 184.

Among 3,318 females who were housewives, 13% were in

part-time, and 17% in full-time, remunerative employment.

Table 2.8 suggests that during the first three triennia there was a slight trend towards higher social class, reversed in the last two triennia. The development of the Emergency Clinic may be a factor in this reversal.

6. Marital Status (Table 2.9)

The proportion of single males has increased over three triennia, from 40% to 44%. This is consistent with the increased proportion of younger patients (Table 2.3).

7. Patients who asked to be Referred (Table 2.13)

The question asked at the patient's first attendance was of the form "Did you, or your relatives or friends, ask your doctor to send you to a psychiatrist?" One-quarter of in-patients and nearly a third of out-patients answered yes to this question, proportions the same as in 1958-60. Asking to be referred was commoner in males and most frequent in the middle age-groups.

8. Country of Birth (Tables 2.16, 2.17)

Table 2.16 is a new table. It shows that 84% of patients were born in the United Kingdom, 8% in other Commonwealth Countries and 8% elsewhere. Of the 1,705 patients born outside the U.K.,

nearly one-third were born in Eire and about one-fifth in Europe. Noteworthy is the relatively large number of males born in Asia (which mostly means those born in India and Pakistan). From most foreign countries there were more males than females, but from the group of "other European countries" there were over twice as many female patients as males.

If we make the assumption that the Asian-born patients are all from the Commonwealth and that those born in "other" countries are all foreign born, then we have the comparison shown in Table 2.17. The Maudsley Hospital is in Camberwell but close to the adjacent borough of Lambeth. Compared with London as a whole, Camberwell has a smaller proportion of persons born in the overseas commonwealth countries whereas Lambeth (which includes Brixton) has a higher proportion. The table suggests that persons born in England and Wales are under-represented among the hospital patients. The foreign-born are much over-represented and those from Scotland and N. Ireland and from Eire somewhat over-represented. But it does not seem likely that there is any over-representation of persons born in the commonwealth countries.

Table 2.1 Numbers of adult patients and discharges in five triennia

Status	1949-51	1952-54	1955-57	1958-60	1961-63
Hospital patients In-patients Out-patients	* 2,636 5,151	* 3,353 6,004	9,554 3,580 6,229	10,403 3,947 6,752	11,502 3,948 7,766
Total discharges In-patient discharges Out-patient discharges	8,725 3,245 5,480	3,641 *	10,626 3,942 6,684	11,906 4,477 7,429	13,026 4609 8,417

^{*}Figure not extracted.

Table 2.2 In-patient discharges, by hospital and year

		Maudsley			Bethlem	
Year	Male	Female	Total	Male	Female	Total
1961 1962 1963	380 379 354	418 443 410	798 822 764	245 283 279	499 451 468	744 734 747
1961-63	1,113	1,271	2,384	807	1,418	2,225
1958-60	1,062	1,268	2,330	774	1,373	2,147

Table 2.3 Age and sex.—3,948 in-patients and 7,766 out-patients

			IN-I	PATIEN	TS		
Age	Ma No.	les %	Fema No.	ales %	Persons No.	61-63 %	58-60 %
Under 20 20— 25— 35—	172 \ 205 \ 336 354	23 20 21	202 \ 226 } 486 486	19 22 21	374 \ 431 \} 822 840	21 21 21	16 24 22
45— 55— 65— 75 and over	291 200 94 24	17 12 6 1	383 271 155 63	17 12 6 3	674 471 249 87	17 12 6 2	17 12 7 2
All ages	1,676	100	2,272 OUT	100 -PATIE	3,948 NTS	100	100
Under 20 20— 25— 35— 45— 55— 65— 75 and over	317 \ 516 \ 1,155 980 623 276 76 \ 17 \	21 29 25 16 7 2	275 \ 442 \} 1,063 918 590 310 170 \ 38 \}	19 28 24 16 8 5	392 \ 958 \} 2,218 1,898 1,213 586 246 \ 55 \}	20 29 23 16 8 4	17 29 25 15 9 5
All ages	3,960	100	3,806	100	7,766	100	100

Table 2.4 Previous in-patient discharges (before 1961).—3,948 in-patients

Number of previous discharges	Males	Females	Persons		Pe 61-63	ersons, % 58-60	55-57
None	1,413 173 50 23 17	1,913 215 82 31 31	3,326 388 132 54 48	}	84 10 6	86 10 4	82 14 4
Total in-patients	1,676	2,272	3,948		100	100	100

Table 2.5 Previous out-patient discharges (before 1958).—7,766 out-patients

Number of previous discharges		Males	Females	Persons		Pe 61-63	ersons, % 58-60	55-57
None 1 2 3 4 5	•••	3,112 455 209 84 49 27	2,782 582 250 93 46 30	5,894 1,037 459 177 95 57	}	76 13 6 2	71 13 9 4	72 14 8 3
6 or more Total out-patien	ts	3,960	3,806	7,766	J	100	100	100

Table 2.6 Religion.—11,502 hospital patients

				Perso	ns, % of	known	
Religion	Males	Females	61-63	58-60		52-54 ¹	49-51 ¹
Church of							
England	3,345	3,958	65	68	71	73	74
Roman Catholic	998	998	18	16	13	12	11
Nonconformist	294	326	5	6	7	6	7
Jewish	156	185	3	4	4	4	4
Other	253	191	4	3	3	3	4
None	362	192	5	3	2	2 }	
Total known	5,408	5,850	100	100	100	100	100
Not known	131	113	(2)	(2)	(2)	(2)	(2)
Total patients	5,539	5,963					

¹In-patients plus out-patients

Table 2.7 Occupation: proportion in certain occupational groups.—11,502 hospital patients

FEMALES	ter's Patient's tion occupation No. % of known	13 169 7	48	36	45	85		435	449		146		2.425	(16) 266 (11)	2,691
	Supporter's occupation No. % kno	356	148	161	140	409	369	308	131	282	211	305	2,820	452	3,272
	MALES . % of known	10	4	9	4	OI	0I	14	9	14	14	∞	100	(4)	,
	MA No.	530	234	2,929	199	523	525	756	328	746	740	431	5,304	235	5,539
			•	•	:	:	:	:	:	:	:	:	:	:	•
			•	:	:	•	:	:	:	•	:	:		•	:
	dn	•	:	•	:	:	:	:	•	:	:	:	:	:	•
	Occupational Group	:	:	:	ers	tions	surance	:	:	:	•	•	:	:	:
i	ıpation	50	:	:	manag	nunica	and ins	hnical	:	:	:	tions		:	:
	Occı	acturin	, etc.	orating	rs and	d comr	nance	and tec	ice	:	:	occupa		:	:
		Metal manufacturing	Wood, paper, etc.	Building, decorating	Administrators and managers	Iransport and communications	Commerce, finance and insurance	Professional and technical	Personal service	[-	lled	Other coded occupations	:	:	:
		Metal	Wood	Buildi	Admir	Trans	Comm	Profes	Person	Clerical	Unskilled	Other	:	:	•
	Code Nos.	110-279	470-579	580-609	610-629	630-709	7.0-7.59	618-09/	861-888	890-895	930-979		Total known	Not known	Total patients

Table 2.8 Social class of males for five triennia.—Hospital patients

						Male	s, % of k	nown	
	Sc	ocial cl	ass		49-51	52-54	55-57	58-60	61-63
I					5	6	7	7	7
Π					14	14	15	16	15
III					52	53	53	50	50
IV		• • •			13	11	10	10	10
V	• • •	• • •	• • •		16	16	15	17	18
Tota	al knov	vn			100	100	100	100	100
Not	knowi	ı	• • •	• • •	(2)	(3)	(4)	(6)	(6)
Tota	al patie	ents		• • •	3,776	4,411	4,486	4,907	5,539

Table 2.9 Marital status.—11,502 hospital patients

Marital state	No.	ales % of known	Females No. % of known			n A.C. ¹ Females
Single Married:	2,386	44	1,804	30	31.5	28.4
Not separated Separated (non	2,471	45	3,230	55]	
judicial) Separated (judicial)	290 37	} 6	246 69	4 1	64.1	56.0
Divorced	135 106	3 2	170 413	3 7	1.0	1.6 14.0
T-4-11	5,425	100	5,932	100	100.0	100.0
Not known Total patients	114 5,539	(2)	31 5,963	(1)	100.0	100.0

¹Census 1961, London, aged 15 and over

Table 2.10 Number of children born to patients: expressed as the number of ever-married hospital patients who at the time of admission had none, one, or more, children born alive and alive now.—7,167 ever-married hospital patients.

Nine	nhar of	f		With cl			With children alive now		
Number of children			Males % of No. known		Females % of No. known		Males No.	Females No.	
None 1 2 3 4 5 6 7 8 9 10 and 6	 over		736 743 779 354 186 78 42 16 5 6 8	25 25 26 12 6 4	877 1,049 1,093 536 237 124 58 35 11 6 14	21 26 29 12 6 4	758 761 767 339 178 74 38 14 2 9	915 1,089 1,094 503 236 86 49 24 6 3	
Total know Not know Total patie	n	•••	2,953 86 3,039	100 (3)	4,040 88 4,128	100 (2)	2,947 92 3,039	4,021 107 4,128	

Table 2.11 Birth order: i.e., patients' birth order among children born alive to patients' mothers.—11,502 hospital patients

	Birt	h order	•		Males	Females	Persons % of known
Only child		• • •			555	561	11
1		• • •			1,251	1,292	25
2				• • •	1,108	1,208	24
3				• • •	634	714	14
4		• • •		• • •	460	438	9
5					251	333	6
6		• • •			161	191	4
7			• • •		120	152	
8					62	74	> 7
9 or lower	• • •	• • •	• • •	• • •	46	174	J
Total know			• • •		4,648	5,137	100
Not known		• • •	• • •	• • •	891	826	(17)
Total patient	ts	• • •	• • •	• • •	5,539	5,963	11,502

Table 2.12 Twins, etc.—11,502 hospital patients

	Males	Females	Persons	Persons, % of known
Patients with a twin of:				
Same sex	61	84	145)
Opposite sex	41	51	92	2.2
Sex unknown	11	10	21	
Not known if twin	442	395	837	(7)
Parents first cousins:				(,)
Yes	39	40	79	0.7
Not known	1,495	1,545	3,040	(26)
Total patients	5,539	5,963	11,502	

Table 2.13 Asked to be referred: the proportion of patients who asked to be referred to a psychiatrist, by age.—3,948 in-patients and 7,766 out-patients

		In-patients,	% of known	Out-patients	, % of known
Age		Males	Females	Males	Females
Under 25 25— 35— 45— 55— 65 and over	•••	19 31 32 34 21 19	15 24 23 21 26 19	26 33 36 34 28 23	20 26 32 31 30 26
All ages Not known if asked be referred (as of known)		28 (4)	(3)	<i>33</i> (5)	<i>29</i> (5)
Total patients		1,676	2,272	3,960	3,806

Table 2.14 Previous treatment of hospital patients at other psychiatric units or mental hospitals.—11,502 hospital patients

Number of times treated elsewhere	Males	Females	Persons, % 61-63	% of known 58-60
None Once Twice Three or more Yes, but number of times not stated	3,618 879 343 357 126	3,861 1,096 384 317	67 18 7 6	61 25 9 5
Total known Not known Total patients	5,323 216 5,539	5,762 201 5,963	100 (4)	100 (3)

Table 2.15 Relatives treated psychiatrically: numbers of hospital patients whose relatives had had treatment at the hospital or at some other psychiatric unit or mental hospital.—11,502 hospital patients

	Males	Females	Persons, % of known			
	iviales	Temales	61-63	58-60		
Relatives treated Relatives not treated	1,301 3,075	1,767 3,210	27	31		
Not known	1,163	986	(19)	(16)		
Total patients	5,539	5,963				

Table 2.16 Country of birth.—11,502 hospital patients

	Males	Females	Persons, % of known		
			All countr	ies Outside UK.	
• • •	3,985	4,472	3 84		
• • •	187	187	}		
			16	= 100	
• • •	262	228		29	
• • •	112	114		13	
vealth	52	21		4	
				•	
• • •	60	39		6	
• • •				12	
		0,5		12	
	45	33		5	
,				16	
				6	
•••	94	66		9	
• • •	5,077	5,459	100	100	
•••	462	504	(9)		
	5,539	5,963	11,502	1,705	
	vealth sian ia or	3,985 262 112 vealth 52 sian 60 125 ia or 45 85 70 94 5,077 462	3,985 4,472 187 187 262 228 112 114 52 21 60 39 125 69 ia or 45 33 85 193 70 37 94 66 5,077 5,459 462 504	Males Females All countr 3,985 4,472 187 84 187 187 262 228 112 114 52 21 vealth 52 21 viain 60 39 60 39 69 ia or 45 33 85 193 70 37 94 66 5,077 5,459 5,077 5,459 69	

Table 2.17 Country of birth. Comparison of hospital patients with London population¹. Proportions per 1,000

Country of birth	Hospital patients		London A.C.		Camberwell Borough		Lambeth Borough	
	M.	F.	M.	F.	M.	F.	M.	F.
England and Wales Scotland and N. Ire-	785	820	814	829	887	890	819	845
land	37	34	27	24	20	17	27	22
Eire Commonwealth,	52	42	44	44	31	29	45	42
Colonies, etc	68 ²	442	68	51	48	38	81	60
Foreign	58	60	47	52	14	16	28	31
Total known Not known	1,000 (90)	1,000 (90)	1,000 (5)	1,000 (5)	1,000 (5)	1,000 (5)	1,000 (5)	1,000 (5)

¹Census, 1961

²Includes Asian-born

CHAPTER THREE

ADULTS: HOSPITAL DATA

Introduction

This chapter deals with information relating to the number of spells of care given to patients in the out-patient and in-patient departments. The tables are therefore based on the number of discharges during the triennium, each discharge being the end of a spell of care.

1. Referring Agencies (Tables 3.1-3.4)

In the present triennium, a discharge has been associated with only one—the principal—referring agency. However, for the referring agencies considered below, a comparison with previous triennia is satisfactory.

- (a) In-patients (Table 3.1). Referrals from an Emergency (or Observation) Ward, which had been decreasing in number, increased in the present triennium. In four successive triennia since 1952, the numbers have been 722, 690, 640 and 644, representing proportions of 20, 18, 14 and 14%. Referrals from mental hospitals have also increased, the numbers over four triennia being 63, 54, 42 and 112.
- (b) Out-patients (Tables 3.2-3.4). Referrals from voluntary organizations have increased (from 28 in 1958-60 to 193 now). This probably reflects the activities of such groups as the Samaritan Society. There has also been an increase in the proportion of self-referrals. A table in the previous Report showed that self-referral was commonest in the oldest age-groups. Table 3.4 shows that self-referral is also commonest in Social Class I and II.

If to the 8,417 out-patient discharges (Table 3.3) are added the 2,558 in-patient cases referred directly from the out-patient department (Table 3.1), we obtain a total of 10,975 cases and the distribution of referring agencies for this total is shown in Table 3.3. Over three triennia, referrals from general hospitals, psychiatric consultants and the domiciliary service have decreased.

Of all out-patient discharges, 18% were self-referrals (that is, the patient came to the out-patient department of his own accord). In the previous report (p. 22) it was shown that the self-referral rate tended to be highest in the oldest age-groups. Table 3.4 now shows that the self-referral rate was not appreciably influenced by social class.

2. Duration of Stay (Tables 3.5-3.7)

Over the past four triennia there has been a steady increase in the proportion of in-patient cases staying less than a month—from 15% to 24%—and a corresponding decrease in the proportion staying more than three months—from 37% to 28% (Table 3.6).

This circumstance is expressed in another way in Table 3.5 which shows the median and average duration of stay over five triennia. During this period the average stay has decreased in each sex by more than a month, and the median stay by more than a fortnight. We may thus say that in the past fifteen years (and indeed in the past decade) there has been a 30% reduction in the duration of inpatient stay.

Table 3.7 confirms the finding of the previous Report, that duration of stay is little influenced by social class but that there is a

slight trend for Social Class IV and V to stay longer.

3. Number of Attendances (Table 3.8)

No clear trend is apparent over five triennia in the number of attendances among-out-patients. Compared with the two previous triennia, however, the proportion of cases during 1961-63 seen more than twenty times has diminished while the proportion seen between five and twenty times has notably increased.

4. Special Investigations (Table 3.9)

The proportion of in-patient cases having X-ray examination has risen steadily, the percentages over five triennia being 23, 31, 40, 43 and 54. Bacteriological tests have also become commoner, the percentages rising from 7 to 17. On the other hand there has been a steady fall in cases having electroencephalography (from 28% to 17%) and basal metabolic rate (from 4% to 1%). Cases having special psychological tests of intelligence have also decreased, precipitously in the present triennium, the percentages over five triennia being 54, 54, 40, 39 and 17. The actual numbers of cases during this time having tests of verbal intelligence were 1,763, 1,956, 1,543, 1,729 and 759. An explanation of this fall may be that such tests are increasingly administered by the medical staff.

5. Special Treatment (Tables 3.10, 3.11)

Table 3.10 compares, for five triennia, the numbers of cases in which special forms of treatment were given. Treatment by special drugs has not been listed as records in different triennia are not comparable. Drugs apart, it is apparent from the table that physical methods of treatment have declined in popularity. Coma insulin therapy was discontinued in 1958 and continuous narcosis is now scarcely given. The number of leucotomy operations has fallen by two-thirds (from 91 to 30) since 1952-54, and for electro-convulsive therapy the present triennium shows for the first time a fall in the number as well as in the proportion of cases. There has been little change in the proportion of cases receiving special psychological methods of treatment and the decline in physical methods can be attributed to an increased use of phenothiazine and anti-depressant drugs.

6. OUTCOME OF TREATMENT (Tables 3.12-3.14)

The course of five triennia reveals a small but steady increase in the proportion of cases discharged as recovered or improved, and a steady decrease in those discharged as unchanged or worse (Table 3.12). Whether this reflects improvements in therapy, or the admission of cases with better prognosis, or a trend in medical optimism, is a subject for debate.

There is no appreciable association between outcome and social class (Table 3.13). Of 44 deaths occurring in the hospital during 1961-63 (Table 3.14), a post mortem examination was made in 36 cases. These deaths include eight suicides, where methods employed were: drugs (2 males, 2 females), coal gas (1 male, 1 female), falling under a train (2 females). The numbers of in-patient deaths by

suicide for the past three triennia have been 7, 7 and 8.

7. DISPOSAL (Tables 3.15, 3.16)

(a) In-patients (Table 3.15). In the present triennium only one disposal was recorded for each discharge. This explains the difference between present and past triennia in the referrals to general practitioners. For other disposals, however, comparison of the triennia is satisfactory. Cases discharged to an Emergency (or Observation) Ward have decreased in number and proportion, the numbers over five successive triennia being 140, 139, 101, 65 and 26. On the other hand, the numbers recommended for admission to a mental hospital or psychiatric unit have been largely unchanged (124, 105, 145, 128, 136), though the proportions have decreased slightly. Taken together, these two sources of disposal account for a decreasing number of cases and it may therefore be said that the need to transfer cases to other psychiatric hospitals, either because of the necessity for long-term care or because of disturbed behaviour together with refusal to stay informally, has diminished.

(b) Out-patients (Table 3.16). The number of cases recommended for referral to an Emergency ward increased sharply. Over four triennia, the referral numbers have been 314, 136, 126 and now 451. This increase probably reflects the working of the Mental Health Act of 1959, by which patients can be admitted informally to

Emergency wards.

8. Lapses in Treatment (Tables 3.17, 3.18)

A slight decrease in the proportion of patients discharged against medical advice (from 14% to 12%) may perhaps be distinguished during five triennia; but in the same period a definite and steady increase (from 19% to 27%) is now evident in the proportion of out-patients lapsing in attendance (Table 3.17). This lapse rate was higher among warded out-patients (i.e., patients who were followed-up in the out-patient department after in-patient discharge) than among other out-patients. As warded out-patients will presumably

have been the more ill and given more attention, a lower lapse rate might be expected, and the observed high rate suggests that doctors are over-cautious in discharging follow-up cases.

Table 3.18 examines lapses by social class. There is, perhaps a slight trend in both in-patients and out-patients for Social Class I and II to have the lowest, and Social Class III the highest, lapse rate.

9. Travelling Time to Hospital (Tables 3.19-3.22)

In the present triennium, out-patients were asked at their first attendance the time it had taken them to travel from home to the hospital. This time was not recorded in 35% of cases. Where it was recorded, the time was less than half-an-hour in 54% and more than an hour in 15% (Table 3.19). A long travelling time was more common in Social Class I and II (Table 3.20), in those attending once only (Table 3.21), and in those who became in-patients (Table 3.22). A short travelling time was associated with referral to an Emergency ward and with a high lapse rate.

Table 3.1 Referring agencies for in-patients.—4,609 in-patient discharges

Referring agency	Male Female -		Totals, % of discharges			
Referring agency	Iviale	remate	61-63	58-60	55-57	
Out-patient department	1,082	1,476	56	58	51	
Day-patient department	18	55	2 \(\)			
Emergency unit	310	354	14	14	17	
Psychiatric department of a						
general hospital	157	276	9	8	7	
Domiciliary service	36	107	3	7	7	
Consultant on the hospital staff	98	108	4	4	4	
Consultant not on the hospital				,	·	
staff	22	21	1	1	2	
Non-psychiatric hospital de-					_	
partment	48	83	3	3	3	
Mental hospital	43	69	2	1	1	
General practitioner	21	51	$\frac{2}{2}$	1	1	
Spontaneous (= self-referral)	20	21	7	1	1	
Other	66	68	3	2	6	
		00				
Total discharges	1,920	2,689	4,609	4,4771	3,942	

¹Although in these triennia a discharge might be associated with more than one referring agency, in fact this rarely occurred.

Referring agencies for out-patients.—8,417 out-patient Table 3.2 discharges

					1946
Referring agency	Male	Female	Totals,	Totals, % of disc	
Referring agency	Whate	remate	61-63	58-60	55-57
General practitioner	2,666	2,761	64	64	65
Spontaneous (=self-referral) ²	788	747	18	19	15
Voluntary organizations	156	37	2	0	0
Probation service, remand		,			
home, court or prison	248	64	4	4	4
Non-psychiatric hospital or					
department	70	86	2	2	3
Psychiatric unit or department	1				
of general hospital	69	90	2	3	3
Domiciliary service	10	30			
Psychiatrist on the hospital staff	19	21			
Psychiatrist not on the hospital					
staff	8	8			
Mental hospital	39	32	}		
Assistance institution	3	9	4	4	4
Emergency unit	17	17			
Labour exchange	8	4			
Children's department	4	19			
Other	192	195	4	4	6
Total discharges	4,297	4,120	100 8,417	100 7,429 ¹	100 6,684 ¹
Total albeital 800 ···	1,201	1,120	0,117	1,-12)	

Table 3.3 Referring agencies to the Out-patient Department

Dafamina a saman	Discharges, % of total			
Referring agency		61-63	58-60	55-57
General practitioners		66.5	66.8	68.0
Spontaneous (= self-referrals)	• • •	16.5	16.5	14.4
General hospitals		4.2	5.2	6.5
Probation services, etc	• • •	3.1	3.0	4.1
Consultants (Bethlem-Maudsley and others)	• • •	0.8	1.5	1.8
Domiciliary service		0.6	1.4	2.2
Mental hospital and Emergency units		1.4	1.3	1.2
Others	• • •	6.9	4.3	1.8
Total, %		100.0	100.0	100.0
Total discharges	• • •	10,975	10,047	8,688

¹Note to Table 3.1 applies ²Including patients who attended at the urging of relatives

Table 3.4 Self-referrals, by social class.—4,297 males and 4,120 female out-patient discharges¹

	Socia	.l c lass		Self-referrals,	% in each S.C
	Boeia	.i Ciass		Male	Female
I				21	19
II				19	22
III	• • •			17	18
IV			• • •	20	17
V	• • •	• • •	• • •	19	17
Tota	ıl of kr	nown S	ocial		
Cl	lass	• • •	• • •	18	17
		referra	ls	778	747
Tota	ıl disch	arges	• • •	4,297	4,120

¹Includes 55 male and 122 female discharges in which the social class was not known.

Table 3.5 Median and average durations of in-patient stay for five triennia.—In-patient discharges

Triennia			N	ledian sta (months)	ıy	Average stay (months)		
			Male	Female	Total	Male	Female	Total
1949-51 1952-54 1955-57 1958-60	•••	•••	2.3 2.4 2.3 2.0	2.6 2.6 2.4 2.2	2.5 2.5 2.3 2.1	3.6 3.7 3.0 2.7	4.0 4.1 3.3 2.9	3.8 3.9 3.2 2.8
1961-63	• • •	•••	1.8	2.0	1.9	2.5	2.8	2.7

Table 3.6 Duration of in-patient stay.—4,609 in-patient discharges

Duration of stay	Male	Male Female -		Totals, %				
	TVIGIO		61-63	58-60	55-57	52-54	49-51	
Less than 1 week 1 week— 2 weeks—	98 92 109	142 137 134	24	20	17	15	18	
3 weeks 2 months	. 195 562 342	219 685 556	} 47	48	46	46	45	
3 months 4 months 6 months 9 months	202 199 81	303 262 142	28	30	35	37	35	
1 year and over	22 18	66 43	1	2	2	2	2	
Total discharges (=100%)	1,920	2,689	4,609	4,477	3,942	3,641	3,245	

Table 3.7 Duration of in-patient stay, by social class.—1,878 male and 2,388 female in-patient discharges of known social class

Duration of stay		Social Class							
(months)	I+II	Males, %	iV+V	Females, % I+II III IV+V					
Less than 1 1— 3 and over	25 49 26	27 46 27	24 47 29	27 45 28	24 46 30	21 47 32			
Total discharges ¹ (=100%)	534	878	466	677	1,289	422			

¹In addition, the social class was unknown in 42 male and 301 female discharges.

Table 3.8 Number of out-patient attendances.—8,417 out-patient discharges

Number o	res. Vicini	Male	Female			Totals, %	THE THE SECTION OF TH	M. Constanting the second of t
	attendances		1 ciliale	61-63	58-60	55-57	52-54	49-51
1 2 3	• • •	1,890 774	1,627 714	42 18	45 19	45 19	42 19	53 17
4	• • •	367 253 614	346 285 676	9 6	9 6	9 5	9 6	8 5
11— 16—	• • •	175 69	197 81	21	13	15		
21— 26— 51—	• • •	46 68 22	54 94 22	} 4	8	7	<i>24</i>	17
76— Over 100	• • •	10	16 8		· · · · · · · · · · · · · · · · · · ·		<u> </u>	
Total discha (=100%)		4,297	4,120	8,417	7,429	6,684	*	5,480

^{*}Figure not extracted. The percentages are based on the total discharges from the Out-patient Department.

Table 3.9 Special investigations on in-patients: number of discharges in which one or more of various investigations were made.—4,609 in-patient discharges.

Investigation	Male	Female	То	otals, %	of disc	charges	
investigation	iviale	remale	61-63	58-60	55-57	52-54	49-51
Laboratory Tests Wasserman or Kahn E.S.R Blood count Glucose tolerance Gastric analysis C.S.F Other Biochemical Bacteriological	1,430 1,632 1,678 14 16 174 904 253	1,976 2,239 2,333 21 19 157 1,278 509	74 84 87 1 1 7 47	86 87 82 1 0 8 75 16	86 85 72 3 1 10 69 12	88 88 58 3 1 11 71 9	85 84 57 3 2 9 61
Other Clinical Tests E.E.G E.C.G Air encephalogram Angiogram Other X-ray B.M.R	405 157 142 33 1,112	76 374 179 123 19 1,390 29	3 17 7 6 1 54 1	7 20 5 43	9 23 5 40 4	5 27 4 31 3	5 28 4 23
Psychological Tests Verbal intelligence Non - verbal intelligence Tests of deterioration Aptitude Other Specialist Opinion	404 354 166 78 199	355 307 141 38 186	17 14 7 3 8	39 35 7 3 11	40 37 6 2 9	54 53 7 2 5	54 54 9 3 4
Specialist Opinion Total discharges	1,920	2,689	4,609	4,477	3,942	4,372	3,245

^{*}Figures not extracted

Table 3.10 Comparison of certain special treatments of in-patients for five triennia showing the number of in-patient discharges in which the treatments were given. Sexes together

Treatment	1.53	1949-51	1952-54	1955-57	1958-60	1961-63
E.C.T Coma insulin therapy Modified insulin Leucotomy Continuous narcosis Drug abreaction Group psychotherapy Hypnosis	•••	 925 198 194 34 35 114 20 25	1,075 210 166 91 33 122 31 21	1,325 166 113 66 19 183 19 32	1,328 54 63 62 6 131 27 16	1,170 none 33 30 7 137 40 59
Total discharges	•••	 3,245	3,641	3,942	4,477	4,609

Table 3.11 Special treatments for in-patients.—4,609 in-patient discharges

Treatment		Male	Female		Γotals, %)
Treatment		Iviaic	1 cmaic	61–63	58-60	55–57
E.C.T		376	794	25	30	34
Modified insulin		8	25	1	1	3
Leucotomy		10	20	1	1	2
Anti-depressant drug		541	939	32		
Phenothiazine drug		728	1,011	38 >	56	40
Other special drugs		480	576	23		
Continuous narcosis	• • •	3	4	0	0	1
Treatment for G.P.I		2	3	0	0	0
Drug abreaction		46	91	3	3	5
Group psychotherapy		18	22	1	1	1
Hypnosis	• • •	37	22	1	0	1
Individual psychotherapy	• • •	197	246	10	15	15
No special treatment	• • •	77	96	4	7	8
Total discharges (=100%)	• • •	1,920	2,689	4,609	4,477	3,942

Table 3.12 Outcome of in patient treatment.—4,609 in-patient discharges

Outcome at	Male	Female		-	Γotals, %	, 0	
discharge	Iviaic	1 Cinaic	61-63	58-60	55-57	52-54	49-51
Recovered Much improved	299 673	503 965	353	52	49	48	47
Improved Slightly improved	340 259	497 324	31	30	30	31	30
Unchanged Worse	304 28	343 30	16	18	21	21	23
Suicide Other death	3 14	5 22					
Total discharges (=100%)	1,920	2,689	3,245	3,641	3,942	4,477	4,609

Table 3.13 Outcome of treatment, by social class.—1,920 male and 2,689 female in-patient discharges

	Social class, % of known							
Outcome at discharge	I+II	Male III	IV+V	I+II	Female III	IV+V		
Recovered Improved Unchanged	52 33 15	51 31 18	51 31 18	55 31 14	54 32 14	59 26 15		
Total of known social class ¹ (=100%)	529	871	460	669	1,279	416		

¹Social class not known in 60 male and 325 female discharges.

Table 3.14 Causes of death in 17 male and 27 female patients

						Males	Females
Primary cerebral neoplasm		• • •	• • •			1	5
Subarachoid haemorrhage	• • •	• • •	• • •			1	1
Subdural haematoma	• • •	• • •	• • •	• • •		1	
Arteriosclerotic or senile deme	ntia	• • •	• • •	• • •		2	1
Carcinoma of bronchus	• • •	• • •	• • •	• • •		3	_
Other malignant neoplasms	• • •	• • •		• • •			3
Pulmonary embolus	• • •	• • •		• • •	• • •	1	1
Bronchopneumonia		• • •		• • •		2	6
Chronic pulmonary fibrosis	• • •	• • •		• • •		1	
Heart disease	• • •			• • •		2	4
Femoral artery thrombosis	• • •	• • •					1
Intestinal volvulus	• • •	• • •	• • •	• • •		1	—
Suicide	• • •			• • •		3	5

Table 3.15 Disposal of in-patients.—4,609 in-patient discharges

Disposal	Male	Female	Totals,	% of d	ischarges
——————————————————————————————————————	Iviaic	1 cmaic	61–63	58–60	55–57
To general practitioner	195	177	. 8	69	65
Further treatment or supervision at the hospital					
Out-patient supervision	1,285	1,883	69	69	61
Out-patient social club	50	91	3	12	7
Out-patient psychotherapy	45	50	2	3	3 2 3
Day-patient	9	42	1 2	1	2
Clinic for epilepsies	48	48	2	2	3
Recommended for treatment or supervision elsewhere					
Out-patient supervision	89	155	5	*	*
Emergency ward	14	12	1	2	3
Mental hospital or other					
psychiatric unit	62	74	3	3	3
Other hospital or unit	29	56	3	<i>3</i> <i>3</i>	4
Residential institution	17	19	}		
Other	77	82	3	3	7
Total discharges	1,920	2,689	4,609	4,4771	3,9421

¹Note to Table 3.1 applies. *Figures not extracted.

Table 3.16 Disposal of out-patients.—8,417 out-patient discharges

Disposal	Male	Female	Totals,	% of di	scharges
Disposar	Male	remaie	61–63	58-60	55–57
To general practitioner	1,646	1,807	40	49	53
Further treatment or supervision at the hospital To surveillance Other	78 32	145 43	$\begin{bmatrix} 3 \\ 1 \end{bmatrix}$	6	5
Recommended for treatment or supervision elsewhere Out-patient department or					
psychiatrist Emergency ward Mental Hospital Other	143 219 275 40	145 232 263 29	3 6 7 1	* 12	* 12
Simple advice given Other No special disposal	255 453 1,156	225 172 959	$\begin{pmatrix} 6 \\ 8 \\ 25 \end{pmatrix}$	37	34
Total discharges	4,297	4,120	8,417	7,429 ¹	6,6841

¹Note to Table 3.1 applies.

Table 3.17 Lapses in treatment.—4,609 in-patient, 8,417 out-patient, and 2,558 warded out-patient discharges

Mode of leaving	Molo		Female		Totals, % of discharges				
9					58-60	55-57	52-54	49-51	
In-patients: Left against advice ¹	217	12	321	12	12	13	14	*	14
Out-patients: Lapsed in attendance	1,216	28	1,024	25	27	21	20		
Warded out-patients Lapsed in attendance	348	32	<i>3</i> 96	27	29	26	* }	21	19

¹Includes absconded and failed to return from leave.

^{*}Figures not extracted.

^{*}Figures not extracted.

Table 3.18 Lapses in treatment, by social class.—4,609 in-patient and 8,417 out-patient discharges

					Lapses,	% of discha	rges of kn	own S.C.
Social class ¹					_	itients Female		atients Female
1+II	• • •	• • •		•••	9	13	26	21
III		• • •	• • •	• • •	13	13	30	26
IV+V	• • •	• • •	• • •	• • •	10	11	28	28

¹Social class not known in 42 male and 301 female in-patient discharges, and 265 male and 675 female out-patient discharges.

Table 3.19 Travelling time from home to hospital.—10,629 patients attending the out-patient department

		Ma	les	Females		
Travellin	ng time	:	No.	% of known	No.	% of known
—15 minutes —30 minutes —45 minutes —1 hour —2 hours —4 hours More than 4 hou	•		591 1,196 478 535 403 68 21	18 36 15 16 12 3	591 1,217 484 527 357 83 15	18 37 15 16 11 3
Total known Not known Total patients	•	• • • •	3,292 1,724 5,016	100 (52)	3,274 1,979 5,253	100 (60)

Table 3.20 Travelling time, by social class.—Per cent of 6,566 patients attending the out-patient department and whose travelling time was known. Sexes together

Social cla	ıss		—15 min.	Travellin —30 min.	g Time —1 hr.	over 1 hr.
$egin{array}{cccccccccccccccccccccccccccccccccccc$	• • •	• • •	15 59 26	17 56 27	24 55 21	29 52 18
Total known S.C. not known	• • •	•••	100 (12)	100 (14)	100 (12)	100 (12)

Table 3.21 Travelling time, by number of attendances.—Per cent of 6,566 patients attending the out-patient department and whose travelling time was known

Ni	ımher	of tim	es seer		—15 mins	Travellin		over 1 hr.
141	moer	OI tilli	ics sect	L	15 111115.	50 mms.	1 111.	0 0 0 1 111:
Once					34	38	39	45
Twice		• • •	• • •	• • •	19	20	18	18
3-4	• • •		• • •		18	17	15	14
5-10		• • •	• • •		20	18	19	16
More t	han 10)	• • •	• • •	9	7	9	7
Total					100	100	100	100

Table 3.22 Travelling time by disposal, etc.—Per cent of 6,566 patients attending the out-patient department and whose travelling time was known

	—15 mins	Travellin —30 mins.		over 1 hr.
Disposal to: In-patient department Emergency ward Day hospital	10.6	10.9	10.1	13.5
	2.1	1.4	1.0	0.7
	1.1	1.2	1.4	0.4
First seen at Emergency Clinic Social case work done Lapsed in attendance	50	48	38	29
	12	11	10	9
	30	27	29	25

CHAPTER FOUR

ADULTS: DIAGNOSTIC DATA

1. Numbers of Patients with Various Diagnoses

(a) In-patients. In the first two Reports, diagnostic tables were based on numbers of discharges, whereas in the last three the tables have been based on the numbers of patients. The discharges in any triennium exceed the number of patients by about 10% and the diagnostic distribution of this 10% of re-discharged cases will be weighted in favour of those diagnoses particularly associated with re-admission. Bearing this in mind, however, a reasonable comparison may still be made between the proportions in various diagnostic categories over the five triennia.

Table 4.2 shows there has been a fairly steady rise in the proportion of psychoses and of character disorders, and a corresponding fall in the proportion of neuroses. From Table 4.4 it can be seen that the rise in psychoses has been due, not to schizophrenia where the proportion has remained constant, but to manic-depressive psychoses where the proportion has increased from 18% to 31%. Indeed, because patients with manic-depressive psychosis tend to be re-admitted, their proportion will be over-represented in the first two triennia, so that the true increase (in terms of patients) is greater than the figures show. This increase in manic-depressive psychoses has not been primarily at the expense of neurotic depression, where the proportion has declined only slightly, but it is associated with and largely (though not fully) balanced by a very marked fall in the proportion of patients diagnosed as anxiety neurosis.

It is interesting and perhaps profitable to speculate on the causes of these trends. That there has been a change in the nature of the illnesses is much less likely than that there has been a change in the diagnostic habits of the doctors. Drugs for the treatment of anxiety, while often effective, have not notably increased in efficacy during the last fifteen years, whereas a great deal of confidence has been engendered in the use of new drugs for the treatment of depression. It does not seem improbable that doctors, possibly from the influence of manufacturers' advertisements, have come to see more of the depressive aspect and less of the anxious aspect in the very common case presenting with a mixture of depression and anxiety. Yet if the mantle of anxiety is passing to depression, it is the psychotic and not the neurotic form of depression which is the beneficiary. This is perhaps to be explained on the ground that cases of depressive illness severe enough to need in-patient treatment are more likely to be diagnosed as psychotic.

Cases of hysteria are being admitted or diagnosed less often, the proportion of in-patient cases having fallen from 6% to 3%

and the actual numbers from 174 to 102. The same is true of obsessional neurosis, the number of cases per triennium falling from 99 to 68.

The proportion and number of in-patients with alcoholism and other drug addiction has increased steadily—from 2% to 5% and from 77 to 185 cases. There has also been an increase in cases of pathological and immature personality, though here the increase appears to date only from 1955-57: in that triennium the number was 173 (5%) while in 1961-63 this had risen to 342 (8%). Of possible reasons for these increases, one is that the faster turn-over of other types of case has led to empty beds in the wards and so to the admission of cases in which the need for in-patient treatment is less

pressing or less obvious.

(b) Out-patients. For out-patients, comparison between the diagnostic tables of the first two and of the last three triennia is less satisfactory than for in-patients. This is so because the tables of the first two triennia were not only based on discharges instead of patients but also included warded out-patient discharges (i.e., in-patient cases followed-up in the out-patient department). The total number of cases was thus between 40% and 50% higher than the number of out-patients. Nevertheless, the proportions of cases in the various diagnostic groups may not be seriously distorted and a comparison of these proportions over five triennia is perhaps worth while.

Table 4.6 shows that the diagnostic trends for out-patients are in many ways the same as for in-patients. Thus the proportion of schizophrenic cases is constant, manic-depressive psychosis and all types of character disorders have increased, and anxiety neurosis has greatly decreased. The rate for hysteria has decreased, the numbers of cases over the past three triennia being 223, 152 and 138; and there has also been a slight fall in the number of cases of obsessional neurosis—from 176 to 161 and 125. Unlike in-patients, however, the frequency of neurotic depression among out-patients has risen (from 15% to 20%) and to an extent at least as great as that for manic-depressive psychosis. Here again it may be supposed that cases of depression not severe enough to need in-patient treatment will be more diagnosed as neurotic.

Tables 4.3 and 4.5 show that among males during 1961-63, one-third had an affective disorder (manic-depressive psychosis, neurotic depression or anxiety) and one-third had a character disorder (pathological and immature personality, alcoholism or drug addiction). Among females, these proportions were one-half

and one-tenth.

2. Miscellaneous Diagnoses (Table 4.7)

Over five successive triennia the number of cases of G.P.I. (dementia paralytica) have been 25, 19, 20, 21 and 14; there is thus no sign of a recrudescence of this disease in the twenty years since

the end of the Second World War. The numbers of cases diagnosed as puerperal psychosis have been 52, 98, 76, 64 and 48; either this psychosis has declined in frequency or its diagnosis in popularity, more probably the latter. Over the past nine years, the number of patients with a primary diagnosis of thyrotoxicosis was 8, with a further 13 in which this was an accessory diagnosis; the number of patients with a primary diagnosis of anxiety state during this period was 3,025.

3. Asked to be Referred (Table 4.10)

Among males, in both in-patients and out-patients, those with character disorders were the most likely to have asked their general practitioners to refer them to a psychiatrist. This was not so in the previous triennium, when for in-patients the rate of those asking to be referred was higher in neurosis, and for out-patients there was little difference among the four diagnostic groups.

4. Lapses in Treatment (Table 4.11)

Among the four diagnostic groups, the highest lapse rate occurs in the character disorders. This excess is particularly marked among in-patients where for both sexes the proportion leaving hospital against medical advice is 60% higher than the average of the other groups.

Table 4.1 Diagnosis in four major groups.—11,502 hospital patients 3,948 in-patients and 7,766 out-patients

D'			Ma	_	Fem	ales	Persons,
Diagnostic	group		No.	%	No.	%	%
Neuroses		•••	1,566 1,737	28 32	2,220 2,444	37 41	33 36
Total	ers, etc. 	• • •	1,734 502 5,539	31 9 100	764 535 5,963	13 9 100	29 9 100
Character disorder Miscellaneous	ers, etc.		796 394 336 150 1,676	47 24 20 9 100	1,263 643 208 158 2,272	56 28 9 7 100	52 26 14 8 100
Neuroses Character disorde Miscellaneous .		•••	833 1,329 1,425 373 3,960	21 34 36 9 100	1,041 1,800 571 394 3,806	28 47 15 10 100	24 40 26 10 100

Table 4.2 Diagnosis in four major groups, by triennia

			Patients	s, % of	total	- 0
Diagnostic group	4	9-51	52–54	55–57	58-60	61–63
In-patients						
Psychoses		45	48	51	54	52
Neuroses		36	34	30	27	26
Character disorders, etc		9	8	9	11	14
Miscellaneous		10	10	10	8	8
Total	•	100	100	100	100	100
Out-patients						
Psychoses		26	27	23	25	24
Neuroses		46	47	49	44	40
Character disorders etc		16	16	19	23	26
Miscellaneous		<i>12</i>	10	9	10	10
Total		100	100	100	100	100

Table 4.3 Diagnosis of in-patients.—3,948 in-patients

I.C.D. Code No.	Diagnosis	Males No. %	Females No. %	Persons %
300 301.0 301.1,301.2 302	Psychoses Schizophrenia Manic and circular Depression	268 16 83 5 340 20	339 <i>15</i> 115 <i>5</i> 675 <i>30</i>	15 5 26
303 304, 306 305, 307 308	Paranoid state Senile, etc Alcoholic, epileptic, etc	19 <i>1</i> 31 2 39 2	17 <i>I</i> 40 <i>2</i> 57 <i>2</i>	1 2 2
309	Other	16 <i>1</i>	20 1	1
310 311 312 313 314 315–317 318	Neuroses Anxiety	$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	2 3 1 2 15 2 1
320 (except 320.6), 321 320.6 322, 323	Character disorders, etc. Pathological and immature personality Sexual deviation Alcoholism and drug ad-	149 35 } 11	137	8
324, 325	diction Childhood behaviour and	133 8	52 2	5
326	mental deficiency Other	$\begin{bmatrix} 12 \\ 7 \end{bmatrix}$ 1	8 8 1	1
353 688	Miscellaneous Epilepsies Puerperal states Other	76 <i>5</i> 74 <i>4</i>	60 3 31 1 67 3	3 1 4
	All diagnoses (=100%)	1,676 100	2,272 100	100

Table 4.4 Diagnosis of in-patients, by triennia

D'	40.51		ts, % of		(1 (2
Diagnosis	49–51	52-54	55–57	58–60	61–63
Schizophrenia	15	15	16	17	15
Manic-depressive psychosis	18	20	24	31	31
Neurotic depression	16	16	13	13	15
Anxiety	8	7	6	4	2
Hysteria	6	5	4	3	3
Obsessional neurosis	3	2	2	2	2
Other neurosis	4	5	5	5	5
Pathological and immature per-					
sonality	6	6	5	7	8
Alcoholism and other drug					
addiction	2	2	3	3	5
Total in-patients	3,2451	3,651 ¹	3,580	3,947	3,948

¹In-patient discharges. See text.

Table 4.5 Diagnosis of out-patients.—7,766 out-patients

I.C.D. Code No.	Diagnosis	Males No. %	Females No. %	Persons %
300 301.0 301.1, 301.2 302	Psychoses Schizophrenia Manic and circular Depression	329 8 66 \ 11 360 \}	269 7 85 18 593 }	8 14
303 304, 306 305, 307, 308 309	Paranoid state Senile, etc Alcoholic, epileptic, etc Other	$ \begin{bmatrix} 25 \\ 19 \\ 24 \end{bmatrix} $ 2	$ \begin{vmatrix} 38 \\ 31 \\ 18 \end{vmatrix} $ $ \begin{vmatrix} 7 \end{vmatrix} $	2
310 311 312 313 314 315-317	Neuroses Anxiety Hysteria Phobic Obsessional Depressive With somatic symptoms Other	349 9 53 1 46 1 67 2 587 15 134 3 93 2	347 9 85 2 94 3 58 1 1,001 26 110 3 105 3	9 2 2 1 20 3 3
320 (except 320.6) 321 320.6 322, 323	Character disorders, etc. Pathological and immature personality Sexual deviation Alcoholism and drug ad-	$ \begin{array}{c} 678 \\ 282 \\ 292 \end{array} $ $ \begin{array}{c} 25 \\ 7 \end{array} $	373 \ 10 12 \ 51 \ 1	17 4
324, 325 326	diction Childhood behaviour and mental deficiency Other	67 <i>1</i> 106 <i>3</i>	33 <i>1</i> 102 <i>3</i>	<i>1 3</i>
353 688	Miscellaneous Epilepsies Puerperal states Others Diagnosis uncertain No psychiatric disorder	71 1 	$ \begin{array}{ccc} 69 & 1 \\ 18 \\ 92 \\ 3 \\ 80 & 2 \\ 135 & 4 \end{array} $	2 2 3 3
	All diagnoses (=100%)	3,960	3,806	7,766

Table 4.6 Diagnosis of out-patients, by triennia

Diagnosis		Patien	ts, % of	total	
	49–51		55–57		61–63
Schizophrenia	8	9	8	8	8
Manic-depressive psychosis	10	11	11	14	14
Neurotic depression	15	16	15	18	20
Anxiety	15	15	<i>17</i>	12	9
Hysteria	6	5	4	2	2
Obsessional neurosis	3	3	3	2	\bar{I}
Other neurosis	7	9	11	7	6
Pathological and immature					_
personality	11	12	15	18	17
Alcoholism and other drug					
addiction	2	2	2	3	4
Total out-patients (=100%)	7,7131	8,4991	6,229	6,752	7,766

¹Out-patient and warded out-patient discharges. See text.

Table 4.7 Miscellaneous diagnoses outside Section V (300-326) of the International Classification of Diseases.—Among 11,502 hospital patients

Code No.	Diagnosis	Males	Females
025	Dementia paralytica	10	4
083	Late effects of acute infectious encepha-		
	titis	9	6
193	Malignant neoplasm of the brain	7	8
223	Benign neoplasm of the brain	3	4
241	Asthma	8	6
252	Thyrotoxicosis	0	2
253	Myxoedema	1	4
289.2	Other metabolic diseases	1	4
353	Epilepsy	147	126
354	Migraine	7	10
355	Other brain diseases (including Hunting-		
	don's chorea)	10	10
649	Pregnancy associated with other conditions		14
688.1	Puerperal psychosis		48
726.2	Torticollis	5	4
780	Symptoms referable to the nervous system	9	6
781.6	Speech disturbance	7	1
, 0 1 1 0	Other diagnoses outside 300–326 ¹	66	78
	Total	290	335

¹No rubric containing more than 5 cases.

Table 4.8 Principal accessory diagnoses.—Among 11,502 hospital patients

Code No.	Diagnosis	Males	Females
300-309 310-318 320 321 322 323 325	Psychiatric disorders Psychoses Neuroses Pathological personality Immature personality Alcoholism Other drug addiction Mental deficiency	89 406 396 117 80 26 59	111 501 268 188 30 34 53
002 241 252 253 260 353 444 502 541 649	Non-psychiatric disorders Pulmonary tuberculosis Asthma Thyrotoxicosis Myxoedema Diabetes Epilepsy Essential hypertension Chronic bronchitis Peptic ulcer Pregnancy associated with other conditions	32 4 6 0 0 10 40 9 13 8	23 2 14 3 9 9 53 9 1 2
	Other	143 1,437 4,102	1,624 4,339
	Total patients	5,539	5,963

Marital status by diagnosis.—11,502 hospital patients Table 4.9

	Males				Females			
Diagnostic group	S	M^1	W	B^{2}	S	M^1	W	\mathbf{B}^2
Psychoses Neuroses Character disorders,	27 26	30 37	45 32	22 30	35 37	35 46	71 20	29 41
etc Miscellaneous	38 9	24 9	17 6	42	18 10	9 10	6 3	23 7
All diagnoses	100	100	100	100	100	100	100	100
Total patients ³	2,386	2,471	106	462	1,804	3,230	413	485

¹Includes only those whose marriages were unbroken by separation.
²Broken marriages: includes separated (judicial and non-judicial) and

³Includes 114 males and 31 females of unknown marital status.

Table 4.10 Asked to be referred, by diagnosis.—Among 3,948 inpatients and 7,766 out-patients

Diagnosis	In-patients, Males	% of known Females	Out-patients, Males	% of known Females
Psychoses Neuroses Character, disorders,	22	22	31	29
	28	22	32	26
etc Miscellaneous	39	22	35	28
	22	13	27	28
All diagnoses Total patients ¹	27	21	32	27
	1,676	2,272	3,960	3,806

Includes 136 in-patients (3%) and 360 out-patients (5%) not known if asked to be referred.

Table 4.11 Lapses in treatment, by diagnosis.—Among 3,948 inpatients, 7,766 out-patients and 2,558 warded out-patients

Diagnosis		atients st advice %		patients attendance % Female	Warded Out- patients Lapsed in at- tendance % Male Female	
Psychoses Neuroses	10 10	10 14	24 31	21 28	31 35	25 29
Character disorders etc. Mis'llaneous	17 9	19 11	32 17	28 15	39 22	44 17
All diagnoses	12	12	28	25	32	27
Total patients	1,671	2,272	3,960	3,806	1,078	1,480

CHAPTER FIVE

CHILDREN

A. THE ADOLESCENT UNIT

BY P. H. CONNELL AND W. WARREN

HISTORY

The Adolescent In-patient Unit started in 1949 at Bethlem Royal Hospital in two wards previously used for acute male patients. One of these, Tyson East 2, a ward of seventeen beds, was opened for adolescent boys in April 1949 and the other, Tyson East 1, a ward of eighteen beds, was opened for adolescent girls in April 1950.

Since the war, a number of adolescents had been admitted to the various adult wards of the Maudsley Hospital and experience of these patients showed the need for a separate adolescent unit. The advent of the joint hospital made this development possible.

The Adolescent Unit at Bethlem Royal Hospital and the Adolescent Unit at St. Ebba's Hospital, both being established in 1949, were the first in-patient units for adolescents to be opened in the United Kingdom.

Since 1949, the unit has been in full use and there is always a waiting list because applications for admission have continued to exceed the number of beds available. The unit was partially closed for some months in 1962 for structural alterations and improvements.

The Adolescent Unit has never had a monopoly of adolescent in-patients in the joint hospital and a small number of patients of adolescent age have since been treated at most times in other wards. There have also been transfers of certain patients to and from adult wards and the Adolescent Unit, as their condition required. Some of the more mature adolescents have been found, for instance, to be more at home amongst the adult patients, and the acutely suicidal adolescents may require the special care provided in appropriate adult wards. In this way, whilst retaining a certain independence, the Adolescent Unit has taken its place in the functioning of the rest of the joint hospital.

THE AIM AND PRACTICE OF THE ADOLESCENT UNIT

The objective of the Adolescent Unit has been to provide a hospital treatment regime suitable for patients of this age group. The patients are too mature to be handled effectively in the Children's In-patient Unit at the Maudsley Hospital, but too immature for it to be desirable for them to be housed in an adult ward. In practice, this has meant the admission of patients most often between thirteen and sixteen years of age.

The sexes live separately in their own wards, meeting for social occasions. They are under the care of consultants and registrar medical staff.

The nursing staff are of both sexes for the boys' ward and female for the girls' ward.

Daily occupational therapy is provided which is designed to meet the particular needs of this age-group and throughout the week activities such as swimming, gymnastics and games take place.

There is a full-time psychiatric social worker who deals with the needs of the in-patients. Her help is constantly required for the social needs of patients and for the arrangements for their disposal after leaving hospital, as well as in the sphere of keeping in close contact with the families who more often than not also require support and help over their relationships with and understanding of the patients.

Treatment by the psychiatric staff includes individual psychotherapy, group psychotherapy, manipulation of the environment, psycho-pharmacological agents or E.C.T., according to the needs of the individual patient.

The adolescents in the unit live as an active group under the care of staff who look after them and this gives rise to a dynamic situation which in itself appears to be of therapeutic importance for many of the youngsters admitted, at a stage in their development when they are quickly maturing. This has been found to be an innate feature of the treatment, the totality of which can hardly be described in measurable terms but which appears to be important alongside the investigations and treatment carried out according to individual needs.

EDUCATION

Continued schooling is important for adolescents while in hospital for all those who can benefit. The London County Council have since 1952 supplied continuous schooling on a generous scale for the unit. A wide range of educational needs have been met, ranging from remedial teaching of those who are educationally retarded to preparation for G.C.E. at 'O' level.

In 1962, a prefabricated school building was erected at the rear of the Unit.

OUT-PATIENTS

The in-patient adolescent unit is supported by out-patient clinics held by the same medical staff responsible for the in-patients. These clinics are held at the Maudsley Hospital and the same P.S.W., responsible for the adolescent in-patients, attends these clinics. In this way, out-patient assessment of patients referred for consideration for admission to the adolescent unit can take place and follow-up of discharged patients can be provided. One of us has recently started an out-patient clinic in the adult department for the assessment, treatment and follow-up of the older adolescents,

including patients who have been discharged from the in-patient unit and whose age leads to hostility at being seen in a children's department.

STATISTICS OF THE ADOLESCENT UNIT, 1961-1963

These statistics are shown as part of Tables 5.2, 5.3, 5.15, 5.17, 5.23, 5.24, 5.27 and 5.28. During 1961-63, the number of cases discharged from the adolescent unit was 180. The median age was thirteen and a half, no patient being under eleven years old. The median duration of stay was 4.3 months, more than a month longer than the younger children's unit at the Maudsley. About a third of the cases were discharged recovered or much improved, a third somewhat improved, and a third unchanged. One in six left hospital against medical advice. Three diagnostic categories accounted for 75% of the cases: psychoneurosis 35%, secondary habit disorder 24%, and psychosis 16%.

B. BRIXTON CHILD GUIDANCE UNIT

This clinic is owned and administered by the London County Council and staffed by doctors from the Bethlem-Maudsley Hospital. The London County Council provides psychiatric social workers, educational psychologists and secretarial staff. In October 1963, Dr. P. H. Connell became medical director in succession to Dr. W. Warren.

The Brixton clinic has a close association with the Children's Department of the Maudsley Hospital. It is one of the placements for trainee doctors doing their six months' course in child psychiatry.

Tables 5.31 and 5.32 show the staffing and some case statistics for each of the years 1961-63.

C. COMMENTARY ON THE TABLES

TABLE 5.1: Numbers of children in five triennia.

The numbers of in-patients increased from 250 in 1949-52 to 323 in 1955-57 but have since remained steady. The numbers of outpatients have declined.

TABLE 5.2: In-patient discharges by hospital and year.

Over five triennia, the figures have fluctuated. The mean number of discharges per year for the nine years 1955-63 have been: Maudsley boys 37, girls 18; Bethlem boys 21, girls 32.

Table 5.3: Age.

The table gives the age at first admission (in-patients) or first attendance (out-patients) during the triennium. The boys tend to be younger than the girls. Among in-patients, the proportion under

11 years old is 40% for boys, 16% for girls. Thus it may be said that among children suffering from serious psychiatric disorder, under the age of 11 boys are twice as common as girls, while over the age of 13 the reverse is true.

Tables 5.5 and 5.6: Social Class of parents.

The social class of parents tends to be higher for in-patient children than for out-patients; the same was true in the previous triennium. Over five triennia, the social class of all children shows a clear trend towards an increasing proportion of the upper social classes at the expense of the lower. Relative to the population of Greater London, the hospital population is now over-represented in the upper classes and under-represented in the lower.

TABLE 5.7: Marital status of mothers.

There has been no appreciable change during five triennia in the proportions of mothers with various marital status.

TABLE 5.8: Mothers' age at marriage.

Over five triennia there has been little change in the distribution of mothers' age at first marriage. The table shows this distribution for 1961-63 and also that of females in the London and South East area married in 1950 (the year 1950 was taken because the mean age of hospital children during 1961-63 was about 12). The comparison, in so far as it is proper, suggests that the patients' mothers were considerably younger at marriage than expected.

TABLE 5.9: Mother's age at patients' birth.

Again, there has been no appreciable change during five triennia. The age-distribution of patients' mothers does not differ appreciably from that of women in the London and South East area having a live-born child in 1950.

TABLE 5.10: Sibship size.

This has been essentially unchanged since 1949. Patients who were the only child in the family comprise about 16% of all patients.

TABLE 5.14: Relatives treated psychiatrically.

The term "relatives" here means first degree relatives together with grandparents, uncles, aunts and first cousins. No trend is apparent over five triennia; the proportion of patients having relatives treated at the joint hospital varied between 10% and 13%.

TABLE 5.16: Referring agencies of out-patients.

The proportion of out-patient cases referred from general practitioners and from Child Guidance Units has increased steadily over five triennia: for general practitioners, from 25% to 42%; for Child Guidance Clinics, from 5% to 12%. Otherwise there are no clear trends.

Table 5.18.—Duration of in-patient stay.

The table suggests that since 1949 the median stay has been increasing for boys, but decreasing for girls.

TABLE 5.20: Number of out-patient attendances.

The proportion of out-patient cases seen once only has increased steadily, from 28% to 38%, over five triennia. The proportion seen between five and twenty times has decreased, but there has been no change in the proportion seen more than twenty times. The considerable proportion of cases seen once only reflects the fact that many cases are referred by other clinics for a second opinion or for consideration of admission to the in-patient wards.

TABLE 5.25: Outcome of treatment.

There has been little change during five triennia in the proportion of in-patients assigned to different categories. The proportion discharged as recovered, improved and unchanged are the same for in-patients and out-patients, but the outcome in these two groups may be assessed on different criteria.

Tables 5.27 and 5.28: Lapses in treatment.

The proportion of out-patients who lapsed in attendance has remained fairly constant during five triennia. This proportion (about 25%) is somewhat higher than for adult out-patients (20%). The proportion of in-patient children discharged against medical advice has been about 10%, somewhat lower than for adults (13%).

In children, the lapse rate for out-patients is the same for each sex. But for in-patients leaving against medical advice, the rate for girls is about one-and-a-half times that for boys.

Table 5.28 shows that for both sexes the out-patient lapse rate diminishes with age.

Table 5.29: Diagnosis.

This table is a new departure, compared with previous triennial reports where the diagnoses were given according to the International Classification of Diseases. It is an attempt to categorize the diagnosis of children in terms more meaningful than the very broad terms of the International Classification. These categories were developed by the late Dr. Kenneth Cameron and based on his Diagnostic Categories in Child Psychiatry (*Brit. J. Med. Psychol.*, 1955, **28**, 67).

The categories are seen to be well spread out, but with a majority of children presenting with disturbed personal relationships or conduct disorders. Educational problems are also common.

TABLE 5.30: Country of birth.

This new table shows the country of birth of childrens' parents. In a quarter of all cases, the country of birth was not stated; among the remainder, 9% of fathers and 6% of mothers were born elsewhere than in the United Kingdom or Eire.

Table 5.1 Numbers of patients and discharges for five triennia (children)

Status		1949-51	1952-54	1955-57	1958-60	1961-63
Hospital patients In-patients Out-patients	• • •	250 961	* 300 846	1,193 323 888	1,153 320 840	1,098 314 784
Total discharges In-patient discharges Out-patient discharges	•••	1,410 ¹ 284 1,126 ¹	1,260 ¹ 313 947 ¹	1,258 345 913	1,181 331 850	1,111 323 788

^{*}Figures not extracted.

Table 5.2 Children's in-patient discharges, by hospital and year

Year				Maudsley	7	Ветньем1			
1 cai		Male	Female	Total	Male	Female	Total		
1961 1962 1963	•••	• • •	26 34 46	6 14 17	32 48 63	31 22 26	45 21 35	76 43 61	
1961–63	• • •	•••	106	37	143	79	101	180	

¹Adolescent unit.

¹Not strictly comparable with later triennia. See text.

Table 5.3 Age (on admission) of in-patients and out-patients

		∆ oe				In-pa	Out-p	Out-patients		
Age (years)						DSLEY F.	BETHLEM ¹ M. F.		M.	F.
Less than 3–4 5–7 8–9	3	•••	•••	•••	0 8 27 26	0 1 7 9			4 23 42 68	2 13 30 16
10 11 12 13	• • •	•••	•••	•••	11 19 9	5 4 3 3	4 13 15	2 11 22	26 41 55 57	23 12 28 39
14 15 16 and ov	er	•••	•••	•••	2 0 0	1 3 0	20 17 8	28 23 12	84 63 40	44 37 37
All ages Median a	ge	• • •	•••		103 9.2	36 10.2	77 13.4	98 13.5	503 11.9	281 12.4

¹Adolescent unit.

Table 5.4 Religious upbringing.—1,098 hospital children

Religion	Boys Gi		Children	Children, % of known			
	Boys	Onis	Cilidicii	61-63	58-60	55-57	
Church of England Roman Catholic Non-conformist Jewish Other None	453 90 39 14 14 7	304 45 24 5 7 9	757 135 63 19 21 16	75 13 6 2 2 2	74 14 6 3 2 1	77 13 5 2 2 1	
Total known Not known Total children :	617 66 683	394 21 415	1,011 87 1,098	100 (9)	100 (8)	100 (6)	

Table 5.5 Social class of parents.—314 in-patient children and 784 out-patient children (sexes together)

	Soc	cial cla	100		In-p	atients	Out-patients		
	500	ciai Cia	133		No.	% of known	No.	% of known	
I			•••		28	10	42	6	
ĪI	• • •	•••	• • •	• • •	56	19	129	18	
$_{ m III}$	• • •		• • •		165	56	379	54	
IV	• • •				20	7	63	9	
V	• • •	• • •	• • •	• • •	24	8	92	13	
Total	l known				293	100	705	100	
	cnown		• • •		21	(7)	79	(11)	
Total	l childre	n			314	()	784	(- /	

Table 5.6 Social class of parents (per cent), for five triennia.—Inpatients and out-patients together, sexes together

Social class			Greater London				
Social Cia	188	49-51	52-54	55-57	58-60	61-63	1951
I and II III IV and V	•••	16 59 25	20 55 25	24 54 22	23 55 22	25 55 20	21.5 54.7 23.8
Total	• • •	100	100	100	100	100	100.0

Table 5.7 Marital status of mother.—1,098 hospital children

Marital statu	Boys	Girls	Children	Children, % of known			
Marital status		Doys	Onis	Cimaren	61-63	58-60	55-57
Single Married:	• • •	17	13	30	3	3	4
Not separated Separated Divorced	•••	570 42 20	335 23 16	905 65 36	84 6 4	85 5 4	87 4 2
Total known Not known	• • •	672 11	404 11	1,076 22	100 (2)	100 (4)	100 (3)
Total children	•••	683	415	1,098			

Table 5.8 Mother's age at marriage.—1,098 hospital children. Sexes together

Age	of moth	1er		Chi	ldren	Landan and
	rst marri			No.	% of known	London and South East Area ¹ 1950, %
Under 20				152	24	22.42
20		• • •		308	50	34.22
25		• • •		112	18	19.8
30		• • •		29	5	8.4
35				13	2	9.1
40	• • • •	• • •	• • •	7	1	6.1
Total known			• • •	621	100	100.0
Not known	• • • •		• • •	455	(73)	100.0
Not married	• • •	• • •		22	(, 2)	
Total children	• • •	•••	• • •	1,098		
Mean age at m	arriage	• • •	• • •	23.3		26.3

¹Registrar General's Statistical Review of England and Wales for the Year 1950. London, 1952. Part II, page 72.
²Age groups "under 21" and "21–24".

Mother's age at patient's birth.—1,098 hospital children Table 5.9

7							
Age		Boys	Girls	Child No.	% of known	London and South East Area ¹ 1950	
Under 20 20 25 30 35 40 45 and over	•••		29 138 187 135 67 30 6	21 90 112 73 45 15	50 228 299 208 112 45 11	5 24 31 22 12 5 1	4.1 25.8 33.3 20.9 12.4 3.1 0.4
Total known Not known Total children	•••	•••	592 91 683	361 54 415	953 145 1,098	100 (15)	100.0

¹Registrar General's Statistical Review of England and Wales for the Year 1950. London, 1952. Part II, page 143.

Table 5.10 Sibship size: number of children (including patient) born alive to mother at time of child's first admission during the triennium.—1,098 hospital children

Sibship size		Boys	Girls	Children	Childre	n, % of	known
		Boys Giris		Cilidren	61-63	58-60	55-57
1		79	74	153	15	17	16
2		207	124	341	33	32	35
3	• • •	157	84	241	23	21	37
4		80	49	129	13	14 Š	
5	• • •	43	22	65	6	7)	
6	• • •	38	9	47	5	<i>4</i> } <i>5</i> }	12
7	• • •	11	11	22	5	5)	
8 and over	• • •	19	11	30	5		
Total known	•••	634	394	1,028	100	100	100
Not known	•••	49	21	70	(7)	(7)	(9)
Total children	•••	683	415	1,098	(,)	(,)	(2)
				1000 DI 1000	n n.t	A Park Addison Co.	11-57 1 1- 1/647

Table 5.11 Birth order by sibship size (i.e. patient's birth order among children born alive to their mothers, by number of mother's live-born children at time of patients' admission).

—683 male and 415 female hospital children.

			Si	bship s	ize				
Birth order of patient	1	2	3	Boys ¹	5	6	7	Total known	Not known
1 2 3 4 5 6 and over	77	104 100	47 63 48	21 14 29 15	6 7 12 13 4	8 8 5 7 8 2	2 2 4 6 4 12	265 194 98 41 16 14	5 1 2
Totals	77	204	158	79	42	38	30	628	8
				Girls ¹					
1 2 4 5 6 and over	73	76 58	30 32 22	14 9 13 12	5 6 4 3 5	3 2 0 3 0 1	2 5 4 4 7	201 109 44 22 9 8	4
Totals	73	134	84	48	23	9	22	393	5

¹Sibship size not known for 47 males and 17 females.

Table 5.12 Twins, etc.—1,098 hospital children

	Boys	Girls	Children	Children,	% of known
	Doys	Onis	Cillidicii	61-63	58-60
Patients with a twin of Same sex Opposite sex Sex unknown Not known if twin	2 6 1 15	5 3 0 2	7 9 1 17	\right\} \(1.6 \) \((1.6) \)	2.1 (1.1)
Parents first cousins Yes Not known	4 60	3 47	7 107	0.7 (11)	0.7 (15)

Table 5.13.—Child cared for by foster-parents or in institution (at time of first admission during the triennium).—314 inpatient children and 784 out-patient children

Cared for	Boys	Girls	Children	Children, %	
- Car va 101	Воуз	Onis	Cilidien	61-63	58-60
In-patients By foster-parents In institution	3 15	2 7	5 22	1.6 7.0	5.9 10.6
Total children	180	134	314		
Out-patients By foster-parents In institution	19 39	9 19	28 58	3.6 7.4	6.7 4.6
Total children	503	281	784		

Table 5.14 Relatives treated psychiatrically (i.e. numbers of children whose relatives had had psychiatric treatment).—1,098 hospital children

Relatives treated		Boys Girls Children		Children, 61-63	% of known 58-60	
At this hospital Yes Not known	•••	77 44	53 26	130 ¹ 70	13 (9)	11 (13)
Elsewhere Yes Not known	• • •	144 94	97 58	241 152	22 (14)	20 (26)
Total children	• • •	683	415	1,098		

¹In 55 of these cases (36 boys, 19 girls), relatives had also been treated psychiatrically elsewhere.

Table 5.15 Referring agencies for in-patient children.—323 inpatient discharges

Poforring agangy	MATT	DSLEY	Dray	Bethlem ¹		Totals, % of discharges		
Referring agency	M.	F.	M.	F.	61-63	58-60	55-57	
Out-patient department	42	17	48	57	51	53	55	
Child guidance unit	33	8	14	20	23	13	14	
Psychiatric unit of general		2	_	~		1.1	1.0	
hospital	6	2	5	5	6	11	10	
Local education authority	9	1			3	4	3	
Non-psychiatric unit of								
general hospital	1	2		3	2	5	3	
General practitioner	1	2	1	1)			
Mental hospital	4	1	2	3	> 7	6	9	
Probation service	2	1		4	j			
Others	8	3	9	8	8	8	6	
Total discharges	106	37	79	101				
<u> </u>								

¹Adolescent unit.

Table 5.16 Referring agencies of out-patient children.—788 outpatient discharges

Referring agency	Boy	Girl	Girl Total	Totals, % of discharges		
		- GIII	Total	61-63	58-60	55-57
General practitioner	198	131	329	42	40	37
Probation service	74	43	117	15	17	17
L.C.C. Children's Committee	40	20	60	8	9	12
Child guidance unit	71	23	94	12	9	8
Non-psychiatric unit of general						
hospital	13	8	21	3	3	6
Parents and spontaneous	28	9	37	5	6	7
Psychiatric unit of general hospital	28	15	43	5	4	4
Local education authority (other				_	•	•
than L.C.C.)	28	14	42	5	4	3
Others	25	20	45	5	8	6
Total discharges	505	283	788	100	100	100

Table 5.17 Duration of in-patient stay (children).—323 in-patient discharges

Duration of stay	Мани	Maudsley		Bethlem ¹		Totals, % of discharges			
	M. F.		M. F.		61-63	58-60	55-57		
Less than 1 week 1 week— 2 weeks— 3 weeks	16 6 2 7	7 1 1 3	3 4 3	4 5 2 6	22	16	20		
1 month— 2 months—	16	6	2 8	11 13	20	24	24		
3 months— 4 months— 9 months—	4 6 19	3 5 2 3	7 16 17	17 14 20	\begin{cases} 22 \\ 26 \end{cases}	} 51	47		
1 year— $1\frac{1}{2}$ years—	4 15	2	10 6 3	5 4 —	} } }	9	9		
Total discharges Median stay (months)	106 3.4	37 2.3	79 5.6	101	100	100	100		

¹Adolescent unit.

Table 5.18 Median duration of in-patient stay, by triennia.—in-patient discharges

		Median duration (months)								
		49-51	52-54	55-57	58-60	61-63				
Boys Girls		3.2	3.9	3.7	4.4	4.9				
Girls	•••	2.4	4.4	3.6	3.8	3.3				
Children	• • •	2.8	4.1	3.7	4.0	3.9				

Table 5.19 Number of out-patient attendances (children).—788 outpatient discharges

Number of attendances			Boy	Girl	Total	Totals, %			
		iances	БОУ	GIII	10141	61-63	58-60	55-57	
1		• • •	169	127	296	38	36	34	
2 3	• • •	• • •	45 24	27 11	72 35	19	19	19	
4 5–10	• • •	• • •	32 78	8 45	40 123				
11–15	• • •	• • •	47	19	66	29	33	34	
16–20 21–25	• • •		26 25	14 6	40 31				
26–50 51–75	• • •	• • •	43 13	23	66 15	[14	12	13	
76–100			3	$\tilde{1}$	4	j			
Over 100	•••								
Total disch	arges	•••	505	283	788	100	100	100	

Table 5.20 Number of out-patient attendances (children), by triennia.
—Out-patient discharges

Nur	nher (of atte	endanc	AC	Out-patients, % of discharges						
Ivuii	1001	or atti	ciidaiic	cs	49-51	52-54	55-57	58-60	61-63		
1					28	32	34	36	38		
2–4	• • •		• • •		22	22	19	19	19		
5-20	• • •				411	33^{1}	34	33	29		
21 and r	nore	• • •	• • •	• • •	9^2	13^{2}	13	12	14		
Total	• • •		•••		100	100	100	100	100		

¹5–18 attendances.

²19 and more attendances.

Table 5.21 Special investigations on in-patient children.—323 inpatient discharges

	Boy	Girl	61-63	To 58-60	tals 55-57
	74	93	167	194	214
	80	99	179	197	185
	90	102	192	177	166
	7	4	11	32	26
	83	66	149	150	95
	39	30	69	76	39
etc.)	9	7	16	24	10
	125	105	232	254	259
					8
- 1			•		234
	1		1	1	2
/	138	107	245	257	261
					258
					11
	-	1	•	_	7
		_			61
• • •	25	8	33	37	74
•••	25	24	49	54	65
• • •	185	138	323	331	345
	etc.)	74 80 90 7 83 39 etc.) 9 125 5 94 1 138 120 4 12 73 25 25	74 93 80 99 90 102 7 4 83 66 39 30 etc.) 9 7 125 105 5 2 94 105 1 120 99 4 — 12 1 73 19 25 8 25 24	74 93 167 80 99 179 90 102 192 7 4 11 83 66 149 39 30 69 etc.) 9 7 16 125 105 232 5 2 7 94 105 199 1 138 107 245 120 99 219 4 — 4 12 1 13 73 19 92 25 8 33 25 24 49	Boy Girl 61-63 58-60 74 93 167 194 80 99 179 197 90 102 192 177 7 4 11 32 83 66 149 150 39 30 69 76 etc.) 9 7 16 24 5 2 7 7 94 105 199 233 1 - 1 1 120 99 219 250 4 - 4 8 12 1 13 10 25 8 33 37 25 24 49 54

Table 5.22 Special investigations on out-patient children.—788 outpatient discharges

			The second second			11. 1. 1. 1. 1. 1. 1.	The lates of
Investig	ation			Boy	Girl	Total	
				Boy	GIII	61-63	58-60
Laboratory tests	• • •	• • •	• • •	16	7	23	35
Clinical tests						7	
Electroencephalogra	am			102	49	151	179
Electrocardiogram				2	1	3	6
X-ray	• • •	• • •	* * •	10	3	13	27
Psychological tests							
Verbal intelligence			• • •	364	195	559	564
Non-verbal intellige	ence	• • •		334	179	513	551
Educational	• • •		• • •	116	54	170	223
Other	• • •	• • •	• • •	25	12	37	42
Specialist opinion	• • •	• • •		1	1	2	16
Total discharges	• • •	• • •	• • •	505	283	788	850

Table 5.23 Special treatments, in-patients and out-patients (children).
—323 in-patient and 788 out-patient discharges

		In-pa		Out-	Out-patient	
Special treatment	Mau M.	Maudsley M. F.		BETHLEM ¹ M. F.		F.
E.C.T	19 32 15		1 7 11 17 18 29 20 10 20 7	7 9 21 42 17 31 23 17 37 7	1 4 48 32 122 34 32 42 14	1 1 23 13 56 9 11 36 13
Total discharges	106	37	79	101	505	283

¹Adolescent unit.

Table 5.24 Outcome of treatment (children).—323 in-patient and 788 out-patient discharges

Outcome on	discl	narge			In-pati		Out-patient, % of known		
Outcome on	disci	narge		Maudsley M. F.		BETHLEM ¹ M. F.		M. F.	
Recovered		• • •		2	11	4	6	9	8
Much improved			• • •	24	11	30	<i>28</i>	19	27
Improved			• • •	28	11	34	24	31	31
Slightly improved				11	22	18	18	12	9
No change			• • •	35	47	14	24	27	24
Worse	• • •		• • •					2	1
Suicide		• • •	• • •			<u> </u>			
Other death	• • •	• • •	• • •	-	_				
Total		• • •		100	100	100	100	100	100
Not stated	• • •	• • •	• • •		—			(63)	(94)
Total discharges	• • •	• • •		106	37	79	101	505	283

¹Adolescent unit.

Table 5.25 Outcome of treatment (children) among in-patients for five triennia, and out-patients for two triennia

Outcome on discharge	In	In-patients, % discharges					Out-patients	
	<u> </u>				61-63	% discharges 58-60 61-63		
Recovered and much improved Improved and slightly im-	37	32	32	31	30	33	30	
Proved No change and worse	40 23	44 24	43 25	48 21	42 28	38 29	42 28	
Discharges:	284	313	345	331	323	500 ¹	456 ¹	

¹In a further 350 and 332 out-patient discharges, the outcome was not stated.

Table 5.26 Disposal of in-patients and out-patients (children).— 323 in-patient and 788 out-patient discharges

Disposal	In-pa	atient	Out-patient	
	Boy	Girl	Boy	Girl
To general practitioner	21	14	198	136
Guidance Unit	64	29	142	62
Further treatment or supervision at the hospital Out-patient supervision Out-patient psychotherapy Clinic for epilepsies	57 1	45 2 2		
Recommended for residential observation or treatment Psychiatric unit or hospital Foster-home, residential school, etc	7 15	7 11	16 9	4 5
Other Outside psychiatrist Other	18 2	26	48 40	30 22
No special disposal ¹	2	2	50	25
Total discharges	185	138	505	283

¹Includes lapses in attendance, simple advice given to self-referrals, etc.

Table 5.27 Lapses in treatment (children).—323 in-patient and 788 out-patient discharges

Mode of leaving	% disch	of arges	Totals, % of discharges				
Mode of leaving	M.	F.	49-51	52-54	55-57	58-60	61-63
In-patients: left against advice Maudsley Bethlem ¹	5 16	8 17	}11	*	*	9	12
Out-patients: lapsed in attendance	26	24	252	20^{2}	22	24	25

¹Adolescent unit.

Table 5.28 Lapses in treatment, by age.—788 out-patient discharges

Age (years)	% l Boy	apsed Girl	
Under 10 10–13 14 and over	28 27 23	33 23 20	
Total discharges	505	283	

Table 5.29 Diagnosis of in-patients and out-patients (children).—314 in-patients and 784 out-patients

Diagnostic group		In-pa	Out-patients			
Diagnostic group	Maul M.	DSLEY F.	Ветн М.	ELEM ¹ F.	M.	F.
Physical handicap or ill-health	2	2	1	1	10	7
Intellectual handicap	13	7	2	1	45	31
Personality variant	3	1	8	7	25	10
Primary habit disorder	10	3	1	4	44	27
Secondary habit disorder	35	7	25	17	228	101
Motor disorder	2		5	1	4	1
Education or work disturbance	3	1	2	9	56	23
Other functional symptoms		1		—	6	6
Psychoneurotic disorder	6	4	25	36	47	39
Psychosomatic disorder			<u> </u>		5	3
Organic brain damage syndrome	7	4		2	9	7
Psychotic disorder	22	6	8	20	24	26
Total patients	103	36	77	98	503	281

¹Adolescent unit.

²Not strictly comparable with later triennia. See text. *Figures not extracted.

Table 5.30 Country of birth of children's parents.—1,098 hospital children

Country of birth	Fa	ther	Mo	ther
	No.	% of known	No.	% of known
England and Wales Scotland and Ireland Eire West Indies West African Commonwealth Australian Commonwealth, Canada Hungary, Czechoslovakia, Poland Other European country	747 21 28 10 2 15 15	86 2 3 2	787 15 24 8 2 2 5	89 2 3 1
Other African country Asia Other	3 5 6 11 3	3	$\begin{bmatrix} 3\\4\\7 \end{bmatrix}$	2
Total known Not known Total children	870 228 1,098	100 (26)	882 216 1,098	100 (25)

Table 5.31 Staffing at Brixton Child Guidance Clinic

Staff at 31st Decemb	er		1961	1962	1963
Number of sessions by: Psychiatrists Clinical assistants Psychiatric social workers	•••	•••	16 30 30	17 30 38	15 30 40

Table 5.32 Case statistics at Brixton Child Guidance Clinic

	1961	1962	1963
New cases seen Cases closed Applications received Waiting list on 31st December Average number of cases under treatment per month	220 233 273 28 158	203 218 278 54	203 161 279 61 152

CHAPTER SIX

DAY-PATIENTS AND DOMICILIARY VISITS

A. DAY PATIENTS

During the triennium 429 patients were treated at the two day-hospitals. The number of discharges was 504, giving a re-discharge rate during the triennium of 17%. Fourteen per cent of the patients had been day-patients in earlier triennia, and 25% had been inpatients. These figures reflect the fact that a considerable proportion of patients at the day-hospital have recurrent or relapsing disorders which need sometimes in-patient, sometimes day-patient, treatment.

The 429 patients in the present triennium represent an 18% decrease in the number of the previous triennium, a decrease which is to be accounted for partly by a somewhat lower rate of referral to the Bethlem day-hospital but more to an increased duration of stay (Table 6.6). This increased duration of stay, together with the increased proportion of unmarried patients (Table 6.4) and the decreased proportion of older patients (Table 6.2), may be accounted for by the increased number and proportion of patients with character disorder and the decreased number with depression. The number of patients with depression fell from 328 in 1958-60 to 251 in 1961-63 and this fall accounts for four-fifths of the overall fall in numbers. Reasons for this fall might be that the waiting-time for admission was too long or that, with the advent of anti-depressant drugs, psychiatrists became more confident in the out-patient management of depressive illness.

Table 6.7 shows that the proportion of patients receiving electroconvulsive therapy was 19%, compared with 35% in the previous triennium. This no doubt reflects the increased reliance on drugs in the treatment of depression.

Table 6.3 shows that the proportion of day-patients in the upper two social classes was twice as high at Bethlem as at Maudsley, a reflection of the nature of the districts surrounding the two dayhospitals.

The proportion of day-patients lapsing in attendance or discharging themselves against advice was 10% for males and 9% for females. The corresponding proportion among in-patients was 12% for each sex.

B. Domiciliary Visits

Table 6.10 shows the numbers of domiciliary visits made by the hospital physicians at the request of general practitioners during five triennia. The numbers increased to a maximum in 1955-57 and have since fallen back almost to the level of 1949-51. The largest factor in this fall has been the development of the emergency clinic at the Maudsley Hospital.

As in previous triennia, females account for 70% of cases seen, compared with just over 50% among hospital cases. Thirty-seven per cent of domiciliary cases are aged 55 or over (Table 6.11) compared with 18% among in-patients and 12% among out-patients. Table 6.13 shows the social class of male domiciliary cases to be higher than that of male hospital cases (30% in the upper classes, compared with 23%) but there is no such difference for females.

The recommended disposal of domiciliary cases over five triennia is shown in Table 6.14. The proportion of cases recommended for further treatment at the Bethlem-Maudsley has declined (from 80% to 39%), while the proportion for home treatment has increased (from 9% to 48%). This probably reflects the changing nature of the cases seen at domiciliary consultations. It seems likely that in the early triennia, general practitioners requested domiciliary visits both for urgent and for non-mobile psychiatric cases; but with the development of the Maudsley emergency clinic, urgent cases are increasingly referred to that clinic, leaving the domiciliary service for house-bound cases.

Table 6.1 Number of day-patient discharges, by hospital and year.— 504 day-patient discharges

Year	Mauds	sley Day-h	ospital	Bethlem Day-hospital			T 1
T Cai	Male	Female	Total	Male	Female	Total	Total Discharges
1961	23	54	77	32	53	85	162
1962	31	52	83	33	62	95	178
1963	35	57	92	32	40	72	164
1961-63	89	163	252	97	155	252	504
1958-60	94	216	310	110	185	295	605
1955-57 ¹	42	231	273	33	70	103	376

¹For Bethlem, only the years 1956-57.

Table 6.2 Age of day-patients.—429 day-patients

Age .			Males	Females	Persons	Persons, %		
						61-63	58-60	55-57
Under 25 25— 35— 45— 55— 65 and over	•••		22 34 43 34 25 6	23 63 88 54 28 9	45 97 131 88 53 15	10 23 31 20 12 4	8 23 24 22 15 8	6 22 22 22 18 20 12
All ages	• • •	•••	164	265	429	100	100	100

Table 6.3 Social class of day-patients.—429 day-patients

	Social class			MAUDSLEY % of known Males Females		BETHLEM % of known Males Females		Hospital % of known Males Female	
I				3	4	7	6	5	5
II		• • •	• • •	10	12	16	28	13	19
\mathbf{III}		• • •		46	60	52	49	49	55
IV	• • •			15	17	7	8	11	13
V	• • •	• • •	•••	26	7	18	9	22	8
Tota	al knov	wn		100	100	100	100	100	100
Not known		(3)	(8)	(1)	(8)	(2)	(8)		
Total patients		81	140	83	125	164	265		

Table 6.4 Marital status of day-patients.—429 day patients

Marital	ctatu	c	Males	Famalas	Persons	Persons, %			
Marital status			Wates	remaies	I CISOIIS	61-63	58-60	55-57	
Single			55	65	120	28	25	20	
Married		• • •	103	177	280	65	68	63	
Divorced		• • •	3	7	10	2	1	7	
Widowed	• • •	• • •	3	16	19	5	6	10	
Total	• • •	•••	164	265	429	100	100	100	

Table 6.5 Referring agencies for day-patients.—504 day-patient discharges

Referring agency		Male	Female	Totals, 61-63	% of di 58-60	scharges 55-57
Out-patient department		144	237	76	81	87
In-patient department	• • •	9	36	9	7	4
Psychiatric department of		1.0	2.1	_	_	.1.
general hospital	• • •	16	21	7	5	*
Domiciliary service		4	7	2	2	4
General practitioner			5	1	1	2
Other	• • •	13	12	5	4	3
Total discharges	• • •	186	318	100	100	100

^{*}Figure included with Other referring agency.

Table 6.6 Duration of stay of day-patients.—504 day-patient discharges

Durati	on of	` stav		Male	Female	Total, %			
	on or	stay		Iviaic	1 cmaic	61-63	58-60		
Less than 1 v	week		• • •	12	16	7			
1 week	• • •	• • •	• • •	8	20	20	22		
2 weeks		• • •		20	22	j			
1 month	• • •	• • •	• • •	34	64	34	42		
2 months	• • •		• • •	34	41	}			
3 months			• • •	31	45	1			
4 months		• • •	• • •	22	41	\$ 27	32		
6 months		• • •	• • •	18	38	16	}		
9 months		• • •	• • •	4	19	}			
1 year				2	8	3	4		
$1\frac{1}{2}$ years and	over	• • •	• • •	1	4	}			
Total dischar				2.6	186				
Median stay	(month	ıs)	• • •	2.9	318	2.7	2.3		

Table 6.7 Special treatments of day-patients.—504 day patient discharges

Special treat	tment		Male	Female	Totals, %		
	intente		Wate	remaie	61-63	58-60	
E.C.T			30	66	19	35	
Modified insulin		• • •		1	0	2	
Special drugs:							
Phenothiazines			69	128	39)	
Anti-depressants			94	179	54	<i>45</i>	
Other		• • •	7	79	17		
Psychotherapy	• • •	• • •	11	58	14	22	
Total discharges		• • •	186	318			

Table 6.8 Outcome of day-patient treatment.—504 day-patient discharges

0.4	: 1		Male	Female	Total, %		
Outcome at d	iscnarge	•	Male	remate	61-63	58-60	
Recovered			15	35 \	39	39	
Much improved Improved	• • •	• • •	62 41	82 <i>∫</i> 59 \	30	31	
Slightly improved	• • •	• • •	16	33 }			
No change Worse	• • •	• • •	42	76) 32 (31	30	
Suicide	• • •	• • •	3	<u> </u>			
Other death	•••			1 J			
Total discharges	• • •	• • •	186	318	100	100	

Table 6.9 Diagnosis of day-patients.—429 day-patients

Diag	gnosis			Males	Females	P	ersons,	%
	5110313			1414103	1 cmaics	61-63	58-60	55-57
Psychosis				77	127	48	51	45
Schizophren	ia		• • •	21	23	10		
Manic-depre			• • •	55	101	37		
Other	• • •	• • •	• • •	1	3	1		
Neuroses	• • •			59	114	40	41	49
Anxiety			• • •	10	14	6		
Hysteria			• • •	2	3	1		
Phobic			• • •	2 4 5	15	4		
Obsessional					5	2		
Depressive	• • •		• • •	31	64	22		
Other	• • •	• • •	• • •	7	13	5		
Character disor	rders,	etc.	• • •	23	23	11	6	4
Pathological			ature					
personalit		• • •	• • •	20	22	10		
Other	• • •	• • •	• • •	3	1	1		
Miscellaneous	•••	•••	• • •	5	1	1	2	2
Total				164	265	100	100	100

¹Includes involutional melancholia.

Table 6.10 Domiciliary visits, for five triennia

				Т	rienniu					
			49-51	52-54	55-57	58-60	61-63	1961	1962	1963
Male Female		• • •	140 305	267 709	374 904	294 706	177 420	63 162	72 124	42 134
Total	• • •		445	976	1,278	1,000	597	225	196	176

Table 6.11 Age of cases seen at domiciliary visits

F	Age			Male	Female	Total, %		
					1 ciliare	61-63	58-60	
nder 25	• • •	• • •		27	42	12	11	
25—	• • •			27	59	15	16	
35—				28	98	21	18	
45—	• • •			28	64	15	17	
55				37	72	18	15	
65—	• • •			21	44	11	13	
and over	• • •	• • •	• • •	9	41	8	11	
ll ages	• • •			177	420	100	100	
ll ages	• • •	• • •	• • • (177				

Table 6.12 Marital status of cases seen at domiciliary visits

Mar	rital sta	atus		No.	ale %	Fen No.	nale %
Single Married Divorced Widowed	•••	•••	•••	68 99 1 9	38 56 6	100 243 6 \ 71 }	24 58 18
All	•••	•••	• • •	177	100	420	100

Table 6.13 Social class of cases seen at domiciliary visits

	Soci	al cl	ass		Male % of known	Female % of known
I II IV V		•••	•••		7 23 46 7 17	8 20 55 9 8
Not k	known known ber of vi	 sits	•••	•••	100 (18) 177	100 (80) 420

Table 6.14 Recommended disposal of cases seen at domiciliary visits

	196	51-63	Totals, %					
Recommended Disposal	Male	Female	49-52	52-54	55-57	58-60	61-63	
In-patient (at B-M) Out-patient (at B-M) Day-patient (at B-M) Emergency ward Mental hospital Home treatment Other	41 20 5 11 11 77 12	107 47 15 23 32 168 28	48 33 5 5 9	38 27 9 3 23	23 17 10 9 41	31 15 3 7 11 27 6	25 11 3 6 7 41 7	
Total	177	420	100	100	100	100	100	

Table 6.15 Diagnosis of cases seen at domiciliary visits

						Totals, %	
Diag	nosis			Males	Females	61-63	58-60
Psychoses				72	182	43	48
Schizophrenia				24	53		
Manic and circul	ar			9	20		
Depressive ¹				22	68		
Paranoid state				1	5		
Senile, etc				11	27		
Alcoholic, etc				1	0		
Other		• • •	• • •	4	9		
Neuroses		• • •		72	177	42	36
Anxiety				20	29		
Hysteria				6	29		
Phobic		• • •		3	11		
Obsessional		• • •		2	1		
Depressive	• • •			36	99		
Others	• • •	• • •	• • •	5	8		
Character Disorders	etc			25	35	10	9
Pathological and		e perso		12	9		
Alcoholism and d	rug addi	ction		6	12		
Others				7	14		
Miscellaneous				8	26	5	7
Puerperal states			• • •	_	8		
Others	•••		•••	8	18		
Total				177	420	100	100

¹Includes involutional melancholia.

CHAPTER SEVEN

SPECIAL CLINICS

A. THE OUT-PATIENT DEPARTMENT AND EMERGENCY CLINIC

1. HISTORICAL REVIEW

BY D. H. BENNETT

When the Maudsley Hospital opened in 1923, the out-patient department consisted of a waiting hall, three consulting rooms and a dispensary. Adult patients were seen on four afternoons each week; men on Monday and Thursday and women on Tuesday and Friday. There was no system of appointments. An average of four new adult patients were seen during each session. The main functions of the department were consultation and the selection of patients for admission. At that time, in-patient treatment was considered more efficacious than continued out-patient treatment and in the first year one-third of the new patients attending were admitted. The staff consisted of two senior doctors who saw all new patients and two other doctors. There was one nurse in charge. She undertook all the clerical work and was responsible for the filing of both out-patients

and in-patients' records.

The opinion sometimes maintained, that patients would be unwilling to attend an out-patient clinic at a mental hospital, was not shared by Dr. Mapother and in fact the number of cases seen steadily increased. It was claimed that the department was, at that time, the largest organisation of its kind in the country. By 1929 the number of consulting rooms had been increased. In the following year the Mental Treatment Act legalised expenditure on out-patient treatment and additions were made to the staff. The majority of outpatient attenders came then, as now, from the neighbourhood of the hospital and South London. To extend the hospital's out-patient services the "Northern Psychiatric Clinics" were established in three hospitals north of the Thames in 1931. Three members of the staff, Dr. (now Professor Sir Aubrey) Lewis, Dr. (now Professor) E. W. Anderson and Dr. L. Minski were given charge of clinics at Mile End, St. Charles's and St. Mary's (Highgate) hospital respectively, and attended there twice weekly.

By 1935 the number of new patients seen at the Maudsley was increasing more slowly. The two senior psychiatrists saw four or five new patients each at every session and this was considered a "moderate case load". The work of the department was changing in other ways. The proportion of new patients who were admitted had fallen from 1 in 3 in 1923 to 1 in 14 in 1935. Fewer patients were referred elsewhere for treatment. The number of patients seen more than six times had also increased. One medical officer provided a supportive clinic and gave short interviews to old patients during

the mornings and afternoons. Selected patients attended for psychotherapy. Psychotherapists were employed on an hourly basis and saw patients from the late afternoon until 10 p.m.

Attendances rose from 1,441 in 1923 and 8,223 in 1930 to 19,100 in 1935. More space was needed. To meet this need the present out-patient department was planned and, after delays due to the economic crisis, was opened in 1936. It included, among other provisions, twelve rooms for psychotherapy, the records office and a registry for the clerical staff. In 1939 at the outbreak of war, the hospital was evacuated to Sutton and Mill Hill. During this period and until the hospital re-opened in 1946, the out-patient department at Denmark Hill continued on a reduced basis and was staffed from Sutton Emergency Hospital. In 1948 the implementation of the National Health Service Act and the amalgamation with Bethlem Royal Hospital were associated with a general increase in the medical staff establishment.

The character of the hospital and the function of the out-patient department began to change with the development of a professorial unit, an increased number of registrars in training and the appointment to the joint hospital of more full-time and part-time consultants with specialised interests. New and old patients of both hospitals were seen by appointment at the Maudsley five days a week in the mornings, in the afternoons and evenings. In addition to the many clinics providing consultation and treatment for all types of patient, a number of clinics met special needs. This was not an entirely new development since before the war there had been clinics for post-encephalitic and G.P.I. patients. Following the war, clinics for ex-servicemen and pension cases with occupational problems were provided for a period.

At the present time special clinics deal with alcoholics, elderly, neuropsychiatric and epileptic patients, the problems of forensic psychiatry, and psychotherapy. The institution of the practice and teaching of out-patient group psychotherapy has been a notable advance.

Since 1939 children have been seen in the separate children's department in increasing numbers.

Clinical assistants had been trained in the out-patient department before 1939. The rotation of registrars in training through the various consultant firms and special units led to advances which could not be measured by the number of patients seen. While some patients were still referred, as before, to supportive clinics, an increasing number of consultants were able through their registrars to maintain a continuing responsibility for the treatment of their own patients. Clinics were more widely used by physicians to follow-up discharged in-patients. At present there are 186 doctor's sessions each week, 28 rooms are used and approximately 3,000 out-patients are in active treatment at any given time. From 1949 to 1963 the number of new out-patients has steadily increased (Table 7.1).

The increased number of available consultant appointments did not meet the need of general practitioners for urgent and immediate consultation. In 1951 this demand was met by the establishment of an Emergency Clinic. Patients can be seen at any time during the day or night seven days a week. Many doctors and patients also make enquiries by telephone. This service is unique in this country. That it meets a real need is shown by the steady increase in day-time attendances (Table 7.2; figures for weekend and night-time attendances are not given).

2. Notes on the Tables (7.3-7.6)

The present triennium is the first in which a punch-card analysis has been made of cases seen at the Emergency Clinic. Table 7.3 shows that, during 1961-63, the number of discharges where the spell of care began with an attendance at the Emergency Clinic was 4,491. This is 34% of all discharges from the hospital. Of these 4,491 discharges, 70% were referred by general practitioners and in 18% the patient attended of his own accord (Table 7.4).

Of new patients attending the hospital during 1961-63, 2,261 were first seen at the Emergency Clinic. This is 27% of all new patients. The distribution by age (Table 7.5) and by diagnosis (Table 7.6) of these cases is similar to that of out-patients as a whole.

3. OUT-PATIENT WAITING TIMES

In the Report for 1958-1960 data were presented on in-patient waiting times (page 67 of that report). The present Report gives data on out-patient waiting times—that is, on the interval of time between a general practitioner or other person requesting an out-patient appointment for a patient and the patient's attendance at an out-patient clinic.

Table 7.7 shows the out-patient waiting times for each of the three years of the present triennium. Taking the three years together, we can say that of all patients for whom an out-patient appointment was requested, one-quarter were seen immediately and one-half within three days; two-thirds were seen within a week and three-quarters within a fortnight; 5% of patients waited more than a month to be seen. Compared with 1961, the waiting times show an improvement (i.e., were shorter) in 1962 and 1963.

Of the 7,661 cases in which the waiting time was recorded, 1,411 were first seen at the Emergency Clinic where, from the nature of that clinic, the waiting time was very rarely more than three days. When these 1,411 cases are excluded, we have the figures in the bottom line of Table 7.7—the waiting times for the 6,250 cases first seen at the consultants' clinics. Of these, 56% were seen within a week, 71% within a fortnight, 83% within three weeks, and 94% within a month. Thus the Emergency Clinic substantially affected the proportion of patients seen within a week but did not greatly

affect the proportion of patients seen within three weeks or a month.

Table 7.8 indicates a seasonal fluctuation in waiting time: t e times were shorter in the first and third trimesters, longer in the second and fourth. Considered by month of attendance, the shortest waiting times were in February and August, the longest in July and November. The fluctuations do not appear to be related to the holiday season, but there is a suggestion that bad months (i.e., with long waiting times) were preceded by months with a high attendance rate. Thus the average monthly first attendance rate was 278 (excluding December, when the figure was disproportionately low), but for May, the month before the worst month, it was 325 and for October (before the second worst month) it was 312; on the other hand, it was 303 in July, the month before the second best month. There may also possibly be a causal relation between the seasonal fluctuation of waiting times and the change-over of registrars from one six-month clinical appointment to another in April and October for the longest waiting times occur in the months following these.

B. THE PSYCHIATRIC EMERGENCY UNIT, ST. FRANCIS' HOSPITAL

St. Francis' Hospital, Dulwich, is situated about one mile south-east of the Maudsley Hospital. It is not part of the Bethlem-Maudsley Hospital, but for many years its Observation Ward (called, since 1961, the Emergency Unit) has been staffed by doctors from the Bethlem-Maudsley. Before 1961, patients in this ward were admitted under Section 20 of the Lunacy Act of 1890. Since 1961—that is, since the implementation of the Mental Health Act of 1959—patients are admitted either under one of the sections of the new Act or informally. Case statistics of the Emergency Unit are given in Tables 7.9-7.14.

During 1961-63, there were 3,899 discharges, of whom 60% were admitted under the Mental Health Act and 40% were admitted informally. The proportion of cases with affective disorder was similar to that among the Bethlem-Maudsley in-patients, but the proportion with schizophrenia was twice as high and with personnality disorder four times as high (Table 7.12).

Table 7.14 shows that, of patients admitted informally, nearly one-half were discharged home or to a hostel, and one-half were transferred to a psychiatric hospital. Four per cent of those admitted informally were transferred under the Mental Health Act. Of those admitted on an order under the Mental Health Act, nearly one-third were discharged home or to a hostel and one-fifth were admitted informally to a psychiatric hospital; most of the rest (i.e., some 40% of the total) were transferred to a psychiatric hospital on an order.

C. THE FORENSIC UNIT

An account of this unit was given in the Report for 1958-60 (page 63). Statistics for the present triennium are shown in Table 7.15.

D. THE GUYS-MAUDSLEY NEUROSURGICAL UNIT

The unit was described in the Report for 1952-54 (page 134). Statistics for the present triennium are shown in Tables 7.16-7.18.

Table 7.1 New adult patients referred to out-patient department, 1949-1963

1949-51 ... 5,151 1952-54 ... 6,004 1955-57 ... 6,229 1958-60 ... 6,752 1961-63 ... 7,800 (approx.)

Table 7.2 Emergency clinic attendances (day-time only)

1951	• • •	• • •	395
1952	• • •		666
1953	• • •		863
1958	• • •	• • •	1,830
1959	• • •	•••	1,509
1960	•••		2,274
1963	•••	• • •	2 - 4 -
1705	• • •	• • •	2,010

Table 7.3 Patients first seen at the Emergency Clinic

	No. of	discharges	
Year	Male	Female	Total
1961 1962 1963	659 762 808	641 796 825	1,300 1,558 1,633
Triennium	2,229	2,262	4,491

Table 7.4 Referring agencies to the Emergency Clinic.—4,491 discharges

				То	tal
Referring agency		Male	Female	No.	%
General practitioner	• • •	1,427	1,710	3,137	70
Spontaneous	• • •	463	356	819	18
Voluntary organization	• • •	159	43	202	4
Non-psychiatric hospital	• • •	42	45	87	2
Probation service	• • •	26	7	33	1
Other	• • •	112	101	213	5
Total discharges	• • •	2,229	2,262	4,491	100

Table 7.5 New patients first seen at the Emergency Clinic, by age

A 00		A		Molos	Famalas	Patients		
	Age			Males	Males Females		%	
Under 20				65	72	137 \	19	
20—			• • •	143	161	304		
25—		• • •	• • •	199	174	373 \	30	
30—			• • •	179	146	325 }		
35		• • •		302	252	554	25	
45—		• • •	• • •	198	154	352	16	
55—				86	73	159	7	
65 and over	• • •	• • •	• • •	21	36	57	3	
All ages	• • •	• • •	• • •	1,193	1,068	2,261	100	

Table 7.6 Diagnosis of new cases referred to the Emergency Clinic

Diamaria		Mal	les	Fema	Females		
Diagnosis			No.	%	No.	%	- %
Psychoses	• • •	• • •	291	25	308	29	27
Schizophrenia	• • •	• • •	128		91		10
Manic-depression		• • •	140	12	191	18	15
Other	• • •	• • •	23		26		2
Neuroses		• • •	351	29	492	46	37
Anxiety		• • •	85		61		6
Depression		• • •	215	18	377	35	26
Other	• • •	• • •	51		54		5
Character disorders, e			441	37	172	16	27
Pathological and	imm	ature	_				
1	• • •	• • •	191 \	19	113	11	15
	• • •	• • •	38 \$		25		
Alcoholism and dr	ug a	ddic-					
tion		• • •	181	15	21	2	9 3
Other	• • •	• • •	31		36		3
Miscellaneous	•••	• • •	110	9	96	9	9
Total patients	• • •	• • •	1,193	100	1,068	100	100

Table 7.7 Waiting-times for out-patients. Cumulative percentage of out-patients seen within various times from the date of request for appointment

Year	No wait	3 days	1 week	2 weeks	3 weeks	1 mth.	2 mths.	Total cases of known waiting time	Waiting time not known, as % of known
1961 1962 1963	24 27 27	49 52 53	58 64 64	72 78 76	82 86 86	93 95 95	99 99 99	2,361 2,652 2,638	10 8 7
1961-3 1961-3 (Consultant clinics only ¹)	26 19	51 42	62 56	75 71	85 83	94 94	99 99	7,661 6,250	8 8

¹i.e. omitting the Emergency Clinic. See text.

Table 7.8 Seasonal waiting-time for out-patients, 1961-63.—Cumulative percentage seen within various times

Months	No wait	3 days	1 week	2 weeks	3 weeks	1 mth.	Total cases of known waiting time ¹	Waiting time not known, as % of known
JanMarch	27	52	65	80	89	96	1,720	9
April-June	23	49	60	71	81	92	1,809	9
July-Sept.	25	53	63	75	84	94	1,937	9
OctDec.	24	49	60	74	86	95	1,564	10

¹In 631 cases, the month of attendance could not be extracted (for coding reasons).

Table 7.9 Psychiatric Emergency Unit, St. Francis' Hospital, 1961-63. Status on admission—3,899 discharges

Status on admission ¹	Disch	narges
	No.	%
Informal	1,557	40
Section 25	64 \	2
Section 26	2 }	
Section 29	1,667	43
Section 60	1 7	15
Section 136	606 ∫	
Section 6 (Children's		
Act)	2	
Total	3,899	100

¹Sections (except for Section 6) refer to the Mental Health Act, 1959.

Table 7.10 Psychiatric Emergency Unit, St. Francis' Hospital.— Discharges during 1961-63 by age. Sexes together

A co			Di	scharges
Age	,		No.	%
Under 20			141	4
20		• • •	379	10
25—			1,081	28
35			1,008	26
45			680	16
55			373	10
65 and over	• • •	• • •	237	6
All ages	•••	• • •	3,899	100

Table 7.11 Psychiatric Emergency Unit, St. Francis' Hospital, 1961-63.—Discharges by marital status. 3,899 discharges

Marital status	Disch	arges
Marital Status	No.	%
Single Married ¹ Separated or divorced Widowed	1,898 1,288 438 275	49 33 11 7
Total	3,899	100

¹Excluding those separated.

Table 7.12 Psychiatric Emergency Unit, St. Francis' Hospital, 1961-63. Diagnosis—3,899 discharges

Diagnosis	Disch	arges	
Schizophrenia Affective disorders Psychopathic personality Senile dementia Organic Mental subnormality Neurosis	No. 1,268 1,404 933 80 192 17 5	% 32 36 24 5 2 1	
Total	3,899	100	

Table 7.13 Psychiatric Emergency Unit, St. Francis' Hospital, 1961-63. Referring agency, by status on admission.—From 3,899 discharges

Referring ag	1	tatus on (% of di	Discharges				
	Circy	Informal Sec. 25		Sec. 29 Sec. 136		No. %	
Home or hostel Hospital General practice Police Other	•••	19 77 2 1 1	39 56 — 5	46 51 1 1 1	100	1,100 2,078 44 640 32	28 54 1 16 1
Total, % Total, discharges		100 1,557	100 64	100 1,667	100 606	3,8941	100

¹And five discharges were of other status on admission.

Table 7.14 Psychiatric Emergency Unit St. Francis' Hospital, 1961-1963 Mode of discharge, by status on admission from 3,899 discharges.

Discharged to		1	tatus on (% of di	Discharges			
	Informal	Sec. 25	Sec. 29	Sec. 136	No.	% 	
Psychiatric Hospit Informally Sec. 25 Sec. 26 Home or hostel Other Died	al:	47 3 1 45 4	15 45 2 30 8	23 37 3 3 33 4	18 44 4 31 3	1,237 962 82 1,465 144 15	32 25 2 37 4
Total, % Total, discharges	••••	100 1,557	100 64	100 1,667	100 605	3,8941	100

¹And five discharges were of other status on admission.

Table 7.15 Forensic Unit.—New cases seen and reports made during 1961-63

		1961	1962	1963
New cases seen Maudsley Out-patient Clinic (adults) Maudsley Out-patient Clinic (Children) Brixton prison	•••	89 55 45	105 70 38	110 47 36
Reports made at Remand Homes ¹ Stamford House: To Juvenile Courts For Approved School Classification Mental Health Act	•••	1,242 416 —	1,363 357 14	1,259 368 7

¹No figures available from Cumberlow Lodge Remand Home (girls).

Table 7.16 Numbers of patients and operations.—Neurosurgical Unit

				1961	1962	1963
In-patients from						
Guy's Hospital	• • •		• • •	217	233	191
Maudsley Hospital		• • •		54	61	55
King's College Hospital		• • •		101	81	97
Others	• • •	• • •		104	150	173
Total	• • •	•••	• • •	476	525	516
Out-patients from						
Guy's Hospital	• • •		• • •	857	870	847
Maudsley Hospital		• • •		561	474	461
King's College Hospital	• • •	• • •	• • •		******	
Total	• • •	• • •	• • •	1,418	1,344	1,308
Major operations performed	• • •	• • •	•••	294	314	318
Total operations performed	• • •	• • •	• • •	474	586	576

Table 7.17 Leucotomies.—Operations for Epilepsy

Type of operation					1961	1962	1963
Leucotomy	• • •	• • •	• • •	• • •	19	16	19
Temporal lobe epilepsy	• • •	• • •	• • •	• • •	16	13	11
Other types of epilepsy	• • •	• • •	• • •	• • •	1		1

Table 7.18 Attendances at X-ray Department

Patients from						1961	1962	1963
Neurosurgical unit Bethlem-Maudsley Other hospitals		al	• • •	• • •	• • •	912 1,889 253	999 1,979 219	1,117 1,923 166
Total	• • •	• • •				3,054	3,197	3,206


CHAPTER EIGHT

GENERAL PRACTITIONER REFERRALS

This chapter deals with the number of new adult cases referred by general practitioners to the hospital during 1961-63, and with the location of these practitioners. The location of a practitioner is taken to be the postal address of the place (usually his surgery) from where he made the referral.

Of 6,134 new cases referred to the hospital, 5,117 (84%) were from practitioners in the London postal area and 4,309 (70%) were from practitioners in the South London postal districts (Table 8.1). These proportions are the same as in the previous triennium. The total of new referrals in 1961-63 was 15% higher than that of 1958-60, and the increase was greatest (17%) outside the London area and least (13%) north of the river. However, in the S.W. postal districts of London which are north of the river (see Table 8.3), the number of new cases more than doubled (from 95 to 196) and the proportion of general practitioners referring cases increased from 16% to 29%.

Table 8.3 shows again what was found in the previous two triennia, that "the law of distance" operates and that the nearer a general practitioner is to the Maudsley Hospital, the more likely he is to refer cases there. This is also apparent from the map.






Table 8.1 Numbers of general practitioners and of new cases they referred, by year and location

	1						
Location	1961	ractition 1962	ers 1963	1961	New cas 1962	es 1963	Total of new cases 1961-63
London postal districts: S.E. S.W. W. W.C. E. C. N. M.	296 203 68 16 31 3 27 42	295 227 83 17 21 5 33 50	315 209 85 14 21 4 27 53	927 526 89 25 33 3 37 57	859 592 120 29 27 8 36 66	885 520 127 24 22 6 33 66	2,671 1,638 336 78 82 17 106 189
Counties of England: Essex Hertfordshire Kent Middlesex Surrey Others	18 11 83 19 58 41	14 10 86 26 72 46	24 15 86 28 61 40	19 11 131 20 72 42	14 11 151 32 96 47	24 16 144 29 87 46	51 38 326 81 255 135
Wales Scotland N. Ireland Isle of Wight Isle of Man Channel Islands	3 2 — —	1 1 1 1 1	$\begin{array}{c} 3 \\ 1 \\ \hline 1 \\ \end{array}$	3 2 — —	1 1 1 1	3 1 1 1 —	7 3 1 2 1
Abroad Total	925	991	1,042	2,001	2,094	2,039	6,134

Table 8.2 Numbers of general practitioners in South London postal districts, by number of new cases referred

No of notionts	No. of practitioners					
No. of patients referred	1961	1962	1963			
1	208	211	222			
2	97	99	110			
3	56	71	64			
4	34	44	36			
5	26	24	32			
6	24	29	23			
7	12	16	9			
8	15	10	7			
9	7	4	8			
10	4	3	2			
11-14	8	8	7			
15 and over	5	3	4			

Table 8.3 Number of new cases referred per general practitioner, and proportion of general practitioners referring, by five groups of South London postal districts

Area	Postal districts of area	New Cases referred ¹	G.Ps. referring ¹	G.Ps. on list ²	New cases per G.P. on list	G.Ps referring as % of G.Ps. on list
A	S.E.5, 11, 15, 17, 22, 24,					
n	S.W.9	516	148	156	3.3	.95
В	S.E. 1, 4, 14, 16, 21, 23,	200	107	193	2.0	56
C	S.W.2, 4, 8 S.E.6, 8, 10,	390	107	193	2.0	30
	13, 19, S.W.11, 12, 16, 17	244	98	222	1.1	44
D	S.E.2, 3, 7, 9, 12, 18, 20, 25-27,					
	S.W.13-15, 18-20	221	102	473	0.5	22
E	S.W.1, 3, 5, 6,					
	7, 10 (all north of the river)	65	44	153	0.4	29
	outh London stal districts	1,436	499	1,197	1.2	35

¹Mean of the three years 1961-63.

²Numbers of N.H.S. general practitioners on the list of the London Executive Council and other relevant Councils (1961).

Appendix

I Name and Address of Patient Mr. Mrs. Miss	The Bethlem F and The Maud ADULTS' DI	dsley Hospital EPARTMENT	Number in	
Tel.	5	7 Religion	8 Referred by	1.5
3 Travelling time (Home to O.P.D.)	,		(frames overleaf)	9 No. of previous Discharges
Public transport	Age	C. of E. R.C.	O.P.(I)	
by Private transport	Date of birth	3 4	I.P.	from as as as adult child
4 Maiden name, if married woman	6 Sex	,		
	I M 2 F	5 6 Other None	D.P.	I.P.D.
			O.P.(2)	D.P.D
	8		10-11	12-13
13 Dates of O.P.(1) I.P.	D.P. O.P.(2)	10 Name and	Address of General Practition	oner Telephone No.
Application				
Admission		14		
Discharge		15		
14 Diagnosis	Diagnos Code	itic II Name and	Address of Nearest Relativ (Give Initi	
Principal Disorder				
Accessory Disorder	20-23			
Associated Disorder	24-27	I2 Patient acco		
Other			ompanieu by	Relationship
Disorders	28-31			# ####################################
15 Cause of Death	I I6 P.M. perform	med		
2	I Yes 2 N	***************************************		
		32		
I Retd	work (not if	t and time off h'wife or retd.)	Usual weekly income (less deductions; including overtime)	20 Country of birth
I Retd.	1 Employed	1 Still at work 1	over £25 4 £5-7 7 on N.A	
2 F.T., P.T.	2 Unempl'd	1	£16-25 5 under £5 8 on N.1	4.L.*
H'wife	3 N.K. if empl	Time N.K. 38	£8-15 6 N.K.	N.K.
21 Patient 22 Marital Status asked for referral 1 Single 5 Divorced		e at 24 Duration of present marri	of 25 Age of first 26 Tota age spouse at childr	al number of mother's
I Yes 2 Married 6 Widowed			marriage to patient	alive
2 No 3 Sep. Jud. 7 N.K.			DOFN	[
3 N.K. 4 Sep. Non-Jud. 8 Married m.	oreUnmarried N,K	Unmarried N.	K. Unmarried N.K. now	aliveN.K.
27 Patient's 28 Number of patient's 29 Patien	42	43	44	47
birth order children psychi born alive to	atric unit or hospital	0 Relatives treated at psychiatric unit or hospital		32 Parents were first cousins
mother born aliveN.K. I Yes	If in-patient treat- ment, how many	l Yes	I No 5 Same sex	I Yes
49 2 No	times	2 No	2 Yes, alive 6 Not same	2 No
Now alive N.K. 3 N.K		3 N.K.	3 , Yes, dead 7 5ex N.K.4 N.K. if twin	3 N.K.
N.K. 48 50	N.K. 51	52	r ixix ii twin	53 54 1

FRAME I OUT-PATIENTS (I)	FRAME 2 IN-PATIENTS
40 Previous discharges in 41 Physicians this triennium from	40 Previous discharges in 41 Physicians this triennium from
Hosp., O.P.D.	Hosp, I.P.D.
I None None 55 42 Registrars 56-	56-58 1 None None 55 42 Registrars
2 Formore None	2 formore None
	3 lor more for more
3 lor more lor more 43 Referred from 44 Interim transfer 45 Interim transfer	43 Referred from 144 46 Hospital and Adm. 47 Social Case
to from on	Ward Work
4 D.P.D. 1 O.W. 3 I.P.D. 1.P.D.	2 O P.D. 4 D.P.D. 1 Beth 1 Yes 2 No
4 D.P.D. D.P.D.	2 Maud
46 First seen at 1 47 Social Case 48 Special investigations emergency clinic Work	48 Special Investigations 49 Treatment
I Yes 2 No I Yes 2 No	
49 Treatment 50 Mode of discharge	4 63-64 65-66 50 Discharge 51 Outcome 52 Disposal 53 Duration of
I By Doctor 4 Died	against advice
2 Lapsed 5 Suicide	Yes 2 No
65-66 3 Surveillance	67 68 69-70 71 54a To whom discharged and 54b Next-of-kin (preferably younger
51 Outcome 52 Disposal 53 No. of times see	
	e e e e e e e e e e e e e e e e e e e
68 69.70	55 Date of
54 Follow- up Yes No punching	punching
	EDAME 4
FRAME 3 OUT-PATIENTS (2) 40 Previous discharges in 141 Physicians	FRAME 4 DAY PATIENTS 40 Previous discharges in 1.41 Physicians
40 Previous discharges in this triennium from	FRAME 4 DAY-PATIENTS 40 Previous discharges in this triennium from
40 Previous discharges in this triennium from Hosp. O.P.D.	40 Previous discharges in this triennium from Hosp. D.P.D.
40 Previous discharges in this triennium from	40 Previous discharges in this triennium from Hosp. D.P.D.
40 Previous discharges in this triennium from Hosp. O.P.D. 155 47 Registrars	40 Previous discharges in this triennium from Hosp. D.P.D. 55 42 Registrates
40 Previous discharges in this triennium from Hosp. O.P.D. None None 55 41 Physicians 42 Registrars	40 Previous discharges in this triennium from Hosp. D.P.D. 1 None None 55 42 Registrars
40 Previous discharges in this triennium from Hosp. O.P.D. I None None 2 I or more None 3 I or more I or more 41 Physicians 42 Registrars 42 Registrars 43 Referred from 44 Interim transfer 45 Interim transfer	40 Previous discharges in this triennium from Hosp. D.P.D. I None None 55 2 I or more None
40 Previous discharges in this triennium from Hosp. O.P.D. I None None 2 I or more None 3 I or more I or more 43 Referred from 1 Outside 3 I.P.D. 41 Physicians 42 Registrars 42 Registrars 45 Interim transfer from on	40 Previous discharges in this triennium from Hosp. D.P.D. I None None 2 I or more None 3 I or more I or more 43 Referred from 44 45 46 Hospital
40 Previous discharges in this triennium from Hosp. O.P.D. I None None 2 Lor more None 3 Lor more Lor more 43 Referred from 1 Outside 3 I.P.D. 4 D.P.D. 4 D.P.D. 41 Physicians 42 Registrars 42 Registrars 42 Registrars 45 Interim transfer from on I.P.D. 1 D.W. 3 I.P.D. 1 I.P.D.	40 Previous discharges in this triennium from Hosp. D.P.D. 1 None None 2 Lor more None 3 Lor more Lor more 43 Referred from 44 45 46 Hospital 1 Outside 3 I.P.D. 1 Beth.
40 Previous discharges in this triennium from Hosp. O.P.D. I None None 2 I or more None 3 I or more I or more 1 Outside 3 I.P.D. 4 D.P.D. 59 41 Physicians 42 Registrars 42 Registrars 43 Referred from 1 O.W. 3 I.P.D. 1 D.P.D. 4 D.P.D. 59 59 59 50 51 56 56 56 56 56 56 56 56	40 Previous discharges in this triennium from Hosp. D.P.D. I None None 55 2 I or more None 3 I or more I or more 43 Referred from 44 45 46 Hospital I Outside 3 I.P.D. 2 O.P.D. 59 60 2 Maud.
40 Previous discharges in this triennium from Hosp. O.P.D. I None None 2 I or more None 3 I or more I or more 1 Outside 3 I.P.D. 4 D.P.D. 4 D.P.D. 4 D.P.D. 55 41 Physicians 42 Registrars 42 Registrars 42 Registrars 45 Interim transfer from on I.P.D. 1 O.W. 3 I.P.D. 2 D.P.D. 4 D.P.D. 4 D.P.D. 4 D.P.D. 55.	40 Previous discharges in this triennium from Hosp. D.P.D. 1 None None 2 I or more None 3 I or more I or more 43 Referred from 44 45 46 Hospital 1 Outside 3 I.P.D. 2 O.P.D. 2 Maud.
40 Previous discharges in this triennium from Hosp. O.P.D. I None None 2 I or more None 3 I or more I or more 1 Outside 3 I.P.D. 4 D.P.D. 4 D.P.D. 59 40 Physicians 41 Physicians 41 Physicians 42 Registrars 42 Registrars 45 Interim transfer from on I.P.D. D.P.D. 4 D.P.D. 59 46 First seen at 47 Social Case 48 Special investigations	40 Previous discharges in this triennium from Hosp. D.P.D. I None None 55 2 I or more None 3 I or more I or more 43 Referred from 44 45 46 Hospital I Outside 3 I.P.D. 2 O.P.D. 59 60 2 Maud.
40 Previous discharges in this triennium from Hosp. O.P.D. I None None 2 I or more None 3 I or more I or more 43 Referred from 1 Outside 3 I.P.D. 4 D.P.D. 4 D.P.D. 59 44 Interim transfer from on I.P.D. 4 D.P.D. 59 45 Previous discharges in this trien, serving the serving transfer from on I.P.D. 4 D.P.D. 59 46 First seen at emergency clinic Work I Yes 2 No I Yes 2 No	40 Previous discharges in this triennium from Hosp. D.P.D. I None None 2 I or more None 3 I or more I or more 43 Referred from I Outside 3 I.P.D. 2 O.P.D. 59 48 Special Investigations I Yes 2 No
40 Previous discharges in this triennium from Hosp. O.P.D. I None None 2 I or more None 3 I or more I or more 43 Referred from I Outside 3 I.P.D. 4 D.P.D. 4 D.P.D. 4 D.P.D. 59 48 Special investigations	40 Previous discharges in this triennium from Hosp. D.P.D. 1 None None 55 1 None None 2 I or more None 3 I or more I or more 43 Referred from 44 45 46 Hospital 1 Outside 3 I.P.D. 1 Beth. 2 O.P.D. 59 40 Yes 2 No
40 Previous discharges in this triennium from Hosp. O.P.D. I None None 2 I or more None 3 I or more I or more 42 Registrars 45 Interim transfer from on I.P.D. 4 D.P.D. 4 D.P.D. 4 D.P.D. 59 46 First seen at emergency clinic I Yes 2 No I Yes 2 No 61 41 Physicians 42 Registrars 42 Registrars 45 Interim transfer from on I.P.D. D.P.D. 4 D.P.D. 59 48 Special investigations	40 Previous discharges in this triennium from Hosp. D.P.D. I None None 55 I None None 55 2 I or more None 3 I or more I or more 43 Referred from 44 45 46 Hospital I Outside 3 I.P.D. 1 Beth. 2 O.P.D. 59 60 48 Special Investigations I Yes 2 No 63-64
40 Previous discharges in this triennium from Hosp. O.P.D. I None None 2 I or more None 3 I or more I or more 1 Outside 3 I.P.D. 4 D.P.D. 4 D.P.D. 59 40 First seen at emergency clinic I Yes 2 No I Yes 2 No 61 41 Physicians 42 Registrars 45 Interim transfer from on I.P.D. D.P.D. 4 D.P.D. 59 48 Special investigations 49 Treatment 50 Mode of discharge I By Doctor 4 Died 2 Lapsed 5 Suicide	40 Previous discharges in this triennium from Hosp. D.P.D. I None None 2 I or more None 3 I or more I or more 43 Referred from I Outside 3 I.P.D. 2 O.P.D. 59 48 Special Investigations I Yes 2 No 62 63-64 49 Treatment 41 Physicians 41 Physicians 42 Registrars 42 Registrars 45
40 Previous discharges in this triennium from Hosp. O.P.D. I None None 55 42 Registrars 2 I or more None 3 I or more 1 or more 44 Interim transfer 45 Interim transfer 1 Outside 3 I.P.D. 4 D.P.D. 4 D.P.D. 60 1 D.P.D. 60 1 D.P.D. 60 63 63 64 65 66 66 66 66 66 66	40 Previous discharges in this triennium from Hosp. D.P.D. I None None 55 I None None 55 I None None 3 I or more I or more 43 Referred from 44 45 46 Hospital I Outside 3 I.P.D. I Beth. 2 O.P.D. 59 60 Yaud. 47 Special Case Work 48 Special Investigations I Yes 2 No 62 63-64 49 Treatment 50 Mode of Discharge I By doctor 3 Died 2 Lapsed 4 Suicide
40 Previous discharges in this triennium from Hosp. O.P.D. I None None 2 I or more None 3 I or more I or more 41 Physicians 42 Registrars 42 Registrars 45 Interim transfer from on I.P.D. 4 D.P.D. 4 D.P.D. 4 D.P.D. 59 40 Previous discharges in this triennium from 1 Outside 3 I.P.D. 4 D.P.D. 59 40 P.P.D. 59 40 P.P.D. 50 Mode of discharge 1 By Doctor 4 Died 2 Lapsed 5 Suicide 3 Surveillance	40 Previous discharges in this triennium from Hosp. D.P.D. I None None 55 I None None 55 I Outside 3 I.P.D. 2 O.P.D. 43 Referred from 44 45 46 Hospital I Outside 3 I.P.D. 2 O.P.D. 47 Special Case Work 48 Special Investigations I Yes 2 No 62 63-64 49 Treatment 50 Mode of Discharge I By doctor 3 Died 2 Lapsed 4 Suicide
40 Previous discharges in this triennium from Hosp. O.P.D. I None None 3 I or more I or more 42 Registrars 45 Interim transfer from on I.P.D. 4 D.P.D. 4 D.P.D. 4 D.P.D. 59 46 First seen at emergency clinic I Yes 2 No I Yes 2 No 61 47 Treatment 50 Mode of discharge I By Doctor 4 Died 2 Lapsed 5 Suicide 3 Surveillance	40 Previous discharges in this triennium from Hosp. D.P.D. I None None 55 I None None 55 I None None 3 I or more I or more 43 Referred from 44 45 46 Hospital I Outside 3 I.P.D. I Beth. 2 O.P.D. 59 60 Yaud. 47 Special Case Work 48 Special Investigations I Yes 2 No 62 63-64 49 Treatment 50 Mode of Discharge I By doctor 3 Died 2 Lapsed 4 Suicide
40 Previous discharges in this triennium from Hosp. O.P.D. I None None 3 I or more None 3 I or more I or more 1 Outside 3 I.P.D. 4 D.P.D. 4 D.P.D. 59 44 Interim transfer from on I.P.D. 4 D.P.D. 59 45 Interim transfer from on I.P.D. D.P.D. 59 46 First seen at emergency clinic I Yes 2 No 61 Form on I.P.D. 59 61 50 Mode of discharge I By Doctor 4 Died 2 Lapsed 5 Suicide 3 Surveillance 51 Outcome 52 Disposal	40 Previous discharges in this triennium from Hosp. D.P.D. I None None 55 42 Registrars 2 I or more None 3 I or more I or more 43 Referred from 44 45 46 Hospital I Outside 3 I.P.D. 2 Maud. 61 47 Special Case Work 48 Special Investigations I Yes 2 No 62 63-64 49 Treatment 50 Mode of Discharge 1 By doctor 3 Died 2 Lapsed 4 Suicide 65-66 57 Duration of stay
40 Previous discharges in this triennium from Hosp. O.P.D. I None None 55 42 Registrars 2 I or more None 3 I or more 1 or more 44 Interim transfer 45 Interim transfer 1 Outside 3 I.P.D. 4 D.P.D. 4 D.P.D. 60 1 D.P.D. 60 1 D.P.D. 60 63 63 64 65 66 66 66 66 66 66	40 Previous discharges in this triennium from Hosp. D.P.D. I None None 3 Lor more None 3 Lor more Lor more 43 Referred from I Outside 3 I.P.D. 2 O.P.D. 59 48 Special Investigations I Yes 2 No 62 49 Treatment 50 Mode of Discharge I By doctor 3 Died 2 Lapsed 4 Suicide 65.66 51 Outcome 52 Disposal 53 Duration of stay

WAS LIA STOP

I Name and a	ddress of Patient	and Th	ne Mauds	oyal Hospital sley Hospital EPARTMENT 1963	Number in Register	
		5		7 Religion	8 Referred by (frames overleaf)	9 No. of previous Discharges
3 Name of Sch	ool	Age	6	1 2 C. of E. R.C.	O.P.(I)	from
		Date of birth		3 4	I.P	
4 Secretary Care	Committee	6 Sex		Non-Con. Jewish		O.P.D.
***************************************		1 M.	2 F.	5 6 Other None	O.P.(2)	
Address		*****	7	7 N.K.	O.P. (3)	9-10
13 Dates of	O.P.(1) 1.P.	O.P.(2)	O.P.(3)		ddress of General Practit	
Application						
Admission			-			
Discharge				12		
14 Diagnosis)		Code	II Name and	Address of Nearest Rela	tive Telephone No.
Principal	··· - ···· -	14	· - -			
Accessory		15	·	<u></u>		
Profile				I2 Patient acco	mpanied by	Relationship
; 3.		21 6	26-2			
15 Cause of dear	;h	1	.M. performe	1		
			г	28		
17 Occupation		Retd. N.K.		red for ents' home er-parents' home	19 Country of birth of Mother N.K.	20 Age of mother at patient's birth
2.		F.T., P.T. N.K.	3 in inst	itution	Father N.K.	N.K.
Mother ah		29-33 29-33 22 Mother's age		34	130	Patient's birth order
I Single	5 Divorced	and the state of t	ac iii sc iii ai		ar al	nong children born ive to mother
2 Married	6 Widowed			Born alive	N.K.	
3 Sep. Jud.4 Sep. Non-J.	7 N.K. 8 Married more than once	I to make the second	, , , , , , , , , , , , , , , , , , ,	Alive now	N.K.	
25 Relatives treat	38	Onmarried	N.K.	39 28 Parents first	29 Seen by P.S.W.	42
at B.•M. Hospi	tal at other psychia-		5 Same sex	cousins	27 Seen by 1.3.44.	Yes No
1 Yes			6 Not same se		Name of	
2 No 3 N.K.	1	Yes, deadN.K. if twin	7 Sex N.K.	2 No 3 N.K.	Name of P.S.W	
J IN.N.	43		[a	3 N.K.		

FRAME I OUT-PATIENTS (I)	FRAME 2 IN-PATIENTS				
40 Previous discharges in this triennium from	40 Previous discharges in 41 Physicians this triennium from				
Hosp. O.P.D. 48-50	Hosp. I.P.D. 48-50				
1 None None 47	I None None 47 42 Registrars				
2 Formore None	2 lormore None				
3 For more For more	3 for more for more				
43 Referred from 44 Interim transfer to from on	43 Referred from 44 46 Hospital and Adm. 47 Social Case Ward Work				
1 Outside 3 I.P.D. 3 I.P.D.	1 Outside 1 Beth 1 Yes 2 No				
I.P.D.	2 O.P.D 2 Maud				
46 First seen at 47 Social Case 48 Special Investigations	51 52 53 54 48 Special Investigations 49 Treatment				
emergency clinic Work					
I Yes 2 No I Yes 2 No					
53 54 55.56 49 Treatment 50 Mode of discharge	55-56 57-58 57-58 55-56				
I By Doctor 4 Died	against advice Stay				
2 Lapsed 5 Suicide	Yes 2 No				
51 Outcome 52 Disposal 53 No. of times seen	54a To whom discharged and 54b Next-of-kin (for follow-up) address				
51 Outcome 52 Disposal 53 No. of times seen					
	- Control of the Cont				
60 61-62 63					
54 Follow- up Yes No punching	55 Date of punching				
SPANS 2	FRAME 4 OUT DATIFALTS (2)				
FRAME 3 OUT-PATIENTS (2) 40 Previous discharges in 41 Physicians	FRAME 4 OUT-PATIENTS (3) 40 Previous discharges in 1 41 Physicians				
this triennium from	this triennium from				
Hosp. O.P.D.	Hosp. O.P.D.				
I None None 47 42 Registrars	I None None 47 42 Registrars				
2 lormore None	2 Formore None				
3 for more for more	3 formore formore				
43 Referred from 44 Interim transfer to from on to	43 Referred from 44 Interim transfer 45 Interim transfer to from on				
1 Outside 3 1.P.D. 3 1.P.D.	1 Outside · 3 1.P.D. 3 1.P.D.				
I.P.D.	1.P.D.				
46 First seen at 47 Social Case 48 Special Investigations	46 First seen at 47 Social Case 48 Special Investigations				
emergency clinic Work	emergency clinic Work				
1 Yes 2 No I Yes 2 No	I Yes 2 No I Yes 2 No				
53 54 55.56 49 Treatment 50 Mode of discharge	55.56 49 Treatment 50 Mode of discharge				
. I By Doctor 4 Died	l By Doctor 4 Died				
2 Lapsed 5 Suicide	2 Lapsed S Suicide				
57-58 3 Surveillance 59	57-58 3 Surveillance 59				
51 Outcome 52 Disposal 53 No. of times seen	51 Outcome 52 Disposal 53 No. of times seen				
60 61-62 63	60 61-62				
54 Follow- up Yes No punching	54 Follow- 55 Date of punching				

wallians.

THE BETHLEM ROYAL HOSPITAL AND THE MAUDSLEY HOSPITAL

DOMICILIARY SERVICE

* Encircle as appropriate

Number in Domiciliary Register

					1-5					
NAME OF DOCTOR PAYING VISIT NAME						ADDRESS, TELEPHONE No. SHIP OF NEAREST RELATIVE				
DATE OF VISIT DATE OF										
DATE OF VISIT	APPLICA'	TION	NAME, ADDRESS AND TELEPHONE No. OF PATIENT'S PRIVATE DOCTOR							
8-9		-	-							
AGE AND DATE SE	X	RELIGIO!	N *		MARITAL STATUS *			OCCUPATION		
Age	5. No	on. Con. 4	2. R. C. 3. Jewish 5. None	 Single Widowe Separate Divorce 	ed 4. N. ed 6. Se	arried . K. p. Non-Jud. ngaged		N.K.*	15-18	
PRINCIPAL DIAGNOSIS					DISPOSAL RECOMMENDED					
19-22				2	1. To Wards of Join 2. To Wards of Me 3. To Observation V 4. To Wards of oth	ntal Hospital Ward	5. To C 6. To D 7. Advi	er) D.P.D. Joint Hosp Day Hospital ce to G.P. er disposal	ital	
Approximate distance of home from				Patient's Time of arriving at, and leaving Patient's home						
Bethlem *			•	miles	Arrival					
* Encircle nearest to patient's home Departure										

Printed in England by
WILDING & SON, LTD.
33 CASTLE STREET
SHREWSBURY

